

# Package ‘ZeBook’

June 22, 2026

**Type** Package

**Title** Working with Dynamic Models for Agriculture and Environment

**Version** 1.2

**Maintainer** Francois Brun <francois.brun@acta.asso.fr>

**Description** R package accompanying the book Working with dynamic models for agriculture and environment, by Daniel Wallach (INRAE), David Makowski (INRAE), James W. Jones (U.of Florida), Francois Brun (ACTA), 2019.

**License** LGPL-3

**LazyData** true

**Depends** R (>= 3.0.0)

**Imports** triangle, deSolve

**Suggests**

coda,dlm,maps,mgcv,mnormt,mvtnorm,randomForest,rpart,rpart.plot,sensitivity,sp,spatial,tseries

**LazyDataCompression** xz

**Encoding** UTF-8

**Config/roxygen2/version** 8.0.0

**NeedsCompilation** no

**Author** Francois Brun [aut, cre] (ACTA),  
David Makowski [aut] (INRAE),  
Daniel Wallach [aut] (INRAE),  
James W. Jones [aut] (U.of Florida)

**Repository** CRAN

**Date/Publication** 2026-06-22 17:40:02 UTC

## Contents

AICf . . . . .	3
Bean . . . . .	4
carbonsoil.model . . . . .	5
carbonsoil.update . . . . .	6

carcass.define.param . . . . .	6
carcass.EMI.model . . . . .	7
carcass.EMI.model2 . . . . .	9
carcass.EMI.multi . . . . .	9
carcass.EMI.simule . . . . .	10
carcass.model . . . . .	10
carcass_data . . . . .	12
carrot.weevil.model . . . . .	12
chicks_data . . . . .	13
cotton.model . . . . .	14
epirice.define.param . . . . .	15
epirice.model . . . . .	16
epirice.multi.simule . . . . .	16
epirice.weather . . . . .	17
evaluation.criteria . . . . .	17
exponential.model . . . . .	18
exponential.model.bis . . . . .	19
exponential.model.ie . . . . .	19
goodness.of.fit . . . . .	20
graph_epid . . . . .	21
graph_epid_s . . . . .	21
lactation.calf.model . . . . .	22
lactation.calf.model2 . . . . .	23
lactation.calf.simule . . . . .	24
lactation.define.param . . . . .	24
lactation.machine.model . . . . .	25
lactation.machine.model2 . . . . .	26
magarey.define.param . . . . .	27
magarey.model . . . . .	28
magarey.model2 . . . . .	28
magarey.simule . . . . .	29
maize.data_EuropeEU . . . . .	29
maize.data_MetaModelling . . . . .	30
maize.define.param . . . . .	30
maize.model . . . . .	31
maize.model2 . . . . .	32
maize.muchow.graph . . . . .	33
maize.muchow.model . . . . .	33
maize.multisy . . . . .	35
maize.multisy240 . . . . .	35
maize.RUEtemp . . . . .	36
maize.simule . . . . .	36
maize.simule240 . . . . .	37
maize.simule_multisy240 . . . . .	38
maize.weather . . . . .	39
maize_cir.model . . . . .	39
maize_cir_rue.model . . . . .	40
maize_cir_rue_ear.model . . . . .	41

mm.A.fct . . . . .	42
mm.FAS.fct . . . . .	42
mm.HI.fct . . . . .	43
mm.LN.fct . . . . .	44
param.rtriangle . . . . .	44
param.runif . . . . .	45
population.age.matrix.model . . . . .	45
population.age.model . . . . .	47
population.age.model.ode . . . . .	48
predator.prey.model . . . . .	49
q.arg.fast.runif . . . . .	50
seedweight.data . . . . .	51
seedweight.model . . . . .	51
Sunflower_Phomopsis . . . . .	52
threshold.measures . . . . .	52
verhulst.model . . . . .	53
verhulst.update . . . . .	54
watbal.define.param . . . . .	54
watbal.model . . . . .	55
watbal.model.arid . . . . .	55
watbal.simobsdata . . . . .	56
watbal.update . . . . .	57
watbal.weather . . . . .	57
weather_EuropeEU . . . . .	58
weather_FranceWest . . . . .	59
weather_GNS . . . . .	60
weather_SouthAsia . . . . .	60
weed.define.param . . . . .	61
weed.model . . . . .	62
weed.simule . . . . .	62
weed.update . . . . .	63
WheatYieldGreece . . . . .	63
Wheat_GPC . . . . .	64
zakoks.original.model . . . . .	64
<b>Index</b>	<b>67</b>

AICf

*Calculate AIC, Akaike's Information Criterion***Description**

This function calculate AIC criterion given a vector of observation, a vector of prediction and number of parameter. Note that number of parameters should include variance. AICcomplete is the same calculation of the AIC function of R ( $AIC_{complete} = n \cdot \log(RSS/n) + n + n \cdot \log(2 \cdot \pi) + 2 \cdot p$ , with  $p$  including variance). AICshort is the calculation described in chapter 6 Working with crop model ( $AIC_{short} = n \cdot \log(RSS/n) + 2 \cdot p$ , with  $p$  including variance). difference between AICcomplete and

AICshort is  $AIC_{complete} - AIC_{short} = n + n \cdot \log(2 \cdot \pi)$ . As you use AIC to compare models (with different number of parameters) on a same data (with same  $n$ , number of observation), you can use AICshort or AICcomplete.

### Usage

```
AICf(Yobs, Ypred, npar)
```

### Arguments

Yobs	observed values
Ypred	prediction values from the model
npar	number of parameters (should include variance that count for one supplementary parameter)

### Value

A vector with 'AICcomplete' and 'AICshort'.

### Examples

```
x=c(1,2,3,4,5)
y=c(1.2,1.8,3.5,4.3,5.5)
fit = lm(y~x)
AIC(fit)
AICf(y,predict(fit),3) # 3 parameters : intercept, slope and variance
```

---

Bean

*Bean gene-based models dataset*

---

### Description

Genetic data for the common bean (*Phaseolus vulgaris* L) that was based on a population created by C. E. Vallejos (personal communication; also see Bhakta et al., 2015, 2017) by crossing two widely-differing cultivars of bean (Calima, an Andean type, with Jamapa, a Mesoamerican type).  
 Bean\$marker : Bean Marker Data  
 Bean\$MET : Weather data on 5 Locations  
 Bean\$QTL : QTL Data  
 Bean\$modelpar : Dynamic Model parameters

### Usage

```
Bean
```

### Format

a List including 4 data.frame Bean\$marker, Bean\$MET, Bean\$QTL, Bean\$modelpar.

## Source

C. E. Vallejos (personal communication) and Bhakta et al. (2015, 2017). Bhakta, M. S., V. A. Jones, C. E. Vallejos. 2015. Punctuated distribution of recombination hotspots and demarcation of pericentromeric regions in *Phaseolus vulgaris* L. PLoS ONE 10(1): <https://doi.org/10.1371/journal.pone.0116822>  
 Bhakta, M. S., S. A. Gezan, J. A. Clavijo Michelangeli, M. Carvalho, L. Zhang, J. W. Jones, K. J. Boote, M. J. Correll, J. Beaver, J. M. Osorno, R. Colbert, I. Rao, S. Beebe, A. Gonzalez, J. Ricourte, and C. E.do Vallejos, 2017. A predictive model for time-to-flower in the common bean based on QTL and environmental variables. G3: Genes, Genomes, Genetics 7(12) 3901-3912. <https://doi.org/10.1534/g3.117.300229>.

## Examples

```
# show the maker of JC1 to JC9 values for both parents (JAM and CAL)
# and 5 cRILS (RIJC001 to RIJC005)
Bean$marker[2:8,1:10]
# show the first value of weather data
head(Bean$MET)
# show the value of QTL
Bean$QTL[4:10,1:10]
# show the value of
Bean$modelpar
```

---

carbonsoil.model	<i>The CarbonSoil model - calculate daily values over designated time period</i>
------------------	--

---

## Description

Simple dynamic model of soil carbon content, with a time step of one year. The equations that describe the dynamics of this system are adapted from the Henin-Dupuis model described in Jones et al. (2004). The soil carbon content is represented by a single state variable: the mass of carbon per unit land area in the top 20 cm of soil in a given year (Z, kg.ha<sup>-1</sup>). It is assumed that soil C is known in some year, taken as the initial year. The yearly change in soil C is the difference between input from crop biomass and loss.

## Usage

```
carbonsoil.model(R, b, U, Z1, duration)
```

## Arguments

R	: the fraction of soil carbon content lost per year
b	: the fraction of yearly crop biomass production left in the soil
U	: the amount of C in crop biomass production (constant or time series)
Z1	: initial soil carbon content
duration	: duration of simulation (year)

**Value**

Soil carbon years.

---

carbonsoil.update      *The CarbonSoil model - calculate change in soil carbon for one year*

---

**Description**

Simple dynamic model of soil carbon content, with a time step of one year. The equations that describe the dynamics of this system are adapted from the Henin-Dupuis model described in Jones et al. (2004). The soil carbon content is represented by a single state variable: the mass of carbon per unit land area in the top 20 cm of soil in a given year ( $Z$ , kg.ha<sup>-1</sup>). It is assumed that soil C is known in some year, taken as the initial year. The yearly change in soil C is the difference between input from crop biomass and loss.

**Usage**

carbonsoil.update( $Z_y$ ,  $R$ ,  $b$ ,  $U_y$ )

**Arguments**

$Z_y$	Soil carbon content for year
$R$	the fraction of soil carbon content lost per year
$b$	the fraction of yearly crop biomass production left in the soil
$U_y$	the amount of C in crop biomass production in the given year

**Value**

Soil carbon content for year+1.

---

carcass.define.param      *Define values of the parameters for the Carcass model*

---

**Description**

Define parameters values

**Usage**

carcass.define.param(full = FALSE)

**Arguments**

full                      : if TRUE, return the full description of distribution(default = FALSE)

**Value**

matrix with parameter values (nominal, binf, bsup). A data.frame if full=TRUE

**Examples**

```
carcass.define.param(full=TRUE)
```

---

```
carcass.EMI.model      The Carcass (growth of beef cattle) model with energy as input
```

---

**Description**

**Model description.** This model is proposed by Hoch et. al (2004) to represent the growth of cattle and the relative body composition of different types of animals depending on nutritional conditions. It simulates the dynamics of changes in the composition of the body fat and proteins according to nutrient intake. The system is represented by four state variables: the protein and fat in the carcass (resp. ProtC and LIPC) and other tissues (resp. ProtNC and LipNC) grouped under the name of the fifth district (again, gastrointestinal tract, skin ..). These variables depend on time, the time step used is  $dt = 1$  day. The model is defined by 20 equations, with a total of 18 parameters for the described process.

**Usage**

```
carcass.EMI.model(  
  protcmax,  
  protncmax,  
  alphac,  
  alphanc,  
  gammac,  
  gammanc,  
  lip0,  
  lipc1,  
  lipnc1,  
  beta,  
  delta,  
  amW,  
  b0c,  
  b1c,  
  b0nc,  
  b1nc,  
  c0,  
  c1,  
  energie,  
  PVi,  
  duration  
)
```

**Arguments**

protcmax	amounts of protein in the carcass of the adult animal (kg)
protncmax	amounts of protein in the 5th district of the adult animal (kg)
alphac	maximum protein synthesis rate in the frame (excluding basal metabolism) (j-1)
alphanc	maximum rate of protein synthesis in the 5th district (except basal metabolism) (j-1)
gammac	Maximum rate of protein degradation in the frame (excluding basal metabolism) (j-1)
gammanc	maximum rate of protein degradation in the 5th district (except basal metabolism) (j-1)
lip0	maximum lipid concentration to the theoretical physiological age (percent)
lipc1	increase coefficient of the maximum lipid concentration with the physiological age of the carcass (percent)
lipnc1	increase coefficient of the highest lipid concentration with physiological age area in the 5th (percent)
beta	lipid synthesis rate (j-1)
delta	lipid degradation rate (d-1)
amW	definition
b0c	coefficient of the allometric equation linking mass and lipid-protein carcass
b1c	exponent allometric equation linking mass and defatted protein carcass
b0nc	coefficient of the allometric equation linking mass and lipid-protein 5th district
b1nc	exponent allometric equation linking mass and lipid-protein 5th district
c0	coefficient of the allometric equation between live weight and live weight empty
c1	exponent allometric equation linking body weight and live weight empty
energie	Metabolizable energy available
PVi	initial liveweight
duration	duration of simulation

**Value**

matrix with ProtC,LipC,ProtNC,LipNC,PV

---

carcass.EMI.model2      *The Carcass model function for use with carcass.EMI.simule*

---

**Description**

see carcass.EMI.model for model description.

**Usage**

```
carcass.EMI.model2(param, energie, PVi, duration)
```

**Arguments**

param               : a vector of parameters  
 energie             : Metabolizable energy available  
 PVi                 : initial liveweight  
 duration            : duration of simulation

**Value**

data.frame with PV,ProtC,ProtNC,LipC,LipNC

---

carcass.EMI.multi      *Wrapper function to run Carcass model on several animals with different conditions*

---

**Description**

wrapper function for multisimulation with carcass.EMI.model2

**Usage**

```
carcass.EMI.multi(param, list_individuals, energy, init_condition)
```

**Arguments**

param               : a vector of parameters  
 list\_individuals    : list of individuals  
 energy             : Metabolizable energy available for all individuals  
 init\_condition     : initial condition for all individuals

**Value**

data.frame with id, ration ,duration, day, PV,ProtC,ProtNC,LipC,LipNC

---

carcass.EMI.simule	<i>Wrapper function to the Carcass model for multiple sets of parameter values</i>
--------------------	--

---

**Description**

Wrapper function to the Carcass model for multiple sets of parameter values

**Usage**

```
carcass.EMI.simule(X, energy, PVi, duration)
```

**Arguments**

X	: parameter matrix
energy	: Metabolizable energy available
PVi	: initial liveweight
duration	: duration of simulation

**Value**

data.frame with PV,ProtC,ProtNC,LipC,LipNC

---

carcass.model	<i>The Carcass (growth of beef cattle) model</i>
---------------	--

---

**Description****Model description.**

The model is defined by 20 equations, with a total of 19 parameters for the described process.

**Usage**

```
carcass.model(  
  protcmax,  
  protncmax,  
  alphac,  
  alphanc,  
  gammac,  
  gammanc,  
  lip0,  
  lipc1,  
  lipnc1,  
  beta,
```

```

    delta,
    k,
    b0c,
    b1c,
    b0nc,
    b1nc,
    c0,
    c1,
    cem,
    duration
)

```

### Arguments

protcmax	amounts of protein in the carcass of the adult animal (kg)
protncmax	amounts of protein in the 5th district of the adult animal (kg)
alphac	maximum protein synthesis rate in the frame (excluding basal metabolism) (j-1)
alphanc	maximum rate of protein synthesis in the 5th district (except basal metabolism) (j-1)
gammac	Maximum rate of protein degradation in the frame (excluding basal metabolism) (j-1)
gammanc	maximum rate of protein degradation in the 5th district (except basal metabolism) (j-1)
lip0	maximum lipid concentration to the theoretical physiological age (percent)
lipc1	increase coefficient of the maximum lipid concentration with the physiological age of the carcass (percent)
lipnc1	increase coefficient of the highest lipid concentration with physiological age area in the 5th (percent)
beta	lipid synthesis rate (j-1)
delta	lipid degradation rate (d-1)
k	Parameter coefficient between the half-saturation of the Michaelis-Menten equation of the metabolic weight ( $MJ.kg^{0.75}$ )
b0c	coefficient of the allometric equation linking mass and lipid-protein carcass
b1c	exponent allometric equation linking mass and defatted protein carcass
b0nc	coefficient of the allometric equation linking mass and lipid-protein 5th district
b1nc	exponent allometric equation linking mass and lipid-protein 5th district
c0	coefficient of the allometric equation between live weight and live weight empty
c1	exponent allometric equation linking body weight and live weight empty
cem	parameter for body weight
duration	duration of simulation

### Value

data.frame with ProtC,LipC,ProtNC,LipNC,PV

---

 carcass\_data

*Data of growth of beef cattle for Carcass model*


---

### Description

This dataset is a list of 5 dataframe. list\_individuals : identifiant for each individual energy : Ration (type ration), Individu, time (week), energie (?) init\_condition : Individu, Pvi (initial liveweight) observation\_dynamic :Ration (3 levels "C", "EM", "F"), individu, time, PVobs observation\_slaughter : Ration, individu, time, PVVobs, PVobs, ProtCobs, ProtNCobs, LipCobs, LipNCobs.

### Usage

carcass\_data

### Format

a RangedData instance, 1 row per plot.

### Source

Agabriel, J. (com.pers.)

---

 carrot.weevil.model

*Carrot weevil development model*


---

### Description

**Model description.** Simple model of developpement of carrot weevil.

### Usage

```
carrot.weevil.model(
  tbase = 7,
  ttegs = 130,
  ttlarvae = 256,
  ttprepupae = 114,
  ttpupae = 130,
  ttadultpreovi = 91,
  weather,
  sdate = 1,
  ldate = 360
)
```

**Arguments**

tbase : base temperature  
 tteggs : duration of eggs stage in degre.day  
 ttlarvae : duration of larvae stage in degre.day  
 ttprepupae : duration of prepupae stage in degre.day  
 ttpupae : duration of pupae stage in degre.day  
 ttadultpreovi : duration of adult stage until egg laying in degre.day  
 weather : weather data.frame for one single year  
 sdate : date to begin simulation (day of year) (default 1)  
 ldate : date to end simulation (day of year) (default 360)

**Value**

data.frame with daily state variable

---

chicks_data	<i>Data of growth of chicks</i>
-------------	---------------------------------

---

**Description**

This dataset content dynamic measurements of growth of chicks for different individuals and different strains. The data comes from a selection experiment on chicken initiated by F. Ricard Research Station on Poultry of INRA Nouzilly. The selection focuses on weight at 8 and 36 weeks and allowed to differentiate the following five strains:

strain 1 : X+ (low at 8, but high at 36 weeks)

strain 2 : X+ (high at 8, but low at 36 weeks)

strain 3 : X++ (high weight at both ages)

strain 4 : X- (low weight for both ages)

strain 5 : X0 (control).

This is a sub-sample of 50 females in the last generation of selection with weight data (in g) at 12 different ages to complete measurement (0, 4, 6, 8, 12, 16, 20, 24; 28, 32, 36 and 40 weeks).

**Usage**

carcass\_data

**Format**

a RangedData instance, 1 row : strain ; id\_animal ; time (day) ; liveweight (g).

**Source**

Duval M., Robert-Granie C., Foulley J.-L. (2009) Estimation of heterogeneous variances in non linear mixed models via the SAEM algorithm with applications to growth curves in poultry. *Journal de la Societe Francaise de Statistique*, 150,65-83

Donnet S., Foulley J.-L., Samson A. (2010) Bayesian analysis of growth curves using mixed models defined by stochastic differential equations. *Biometrics* 66, 733-741

Jaffrezic F., Meza C., Foulley J.-L., Lavielle M. (2006) The SAEM algorithm for the analysis of non linear traits in genetic studies. *Genetics, Selection, Evolution*,38, 583-

This dataset was used in a training session Biobayes (France, 2011) training session.

Albert I., Ancelet S., David O., Denis J.B., Makowski D., Parent E., Soubeyrand S. (2012) *Methodes statistiques bayésiennes. Bases theoriques et applications en alimentation, environnement et genetique.* FormaScience. Ecole-chercheurs INRA.

---

 cotton.model

---

*The Cotton model (dynamic for numbers of Cotton fruiting points).*


---

**Description**

**Model description.** TO COMPLETE

**Usage**

```
cotton.model(  
  TESQ,  
  PMAX,  
  AFL,  
  AL,  
  AOP,  
  P1,  
  P2,  
  P3,  
  P4,  
  P5,  
  PF,  
  PSF,  
  TSQ,  
  P,  
  PR,  
  PT,  
  tend  
)
```

**Arguments**

TESQ	: TO COMPLETE
PMAX	: TO COMPLETE
AFL	: TO COMPLETE
AL	: TO COMPLETE
AOP	: TO COMPLETE
P1	: TO COMPLETE
P2	: TO COMPLETE
P3	: TO COMPLETE
P4	: TO COMPLETE
P5	: TO COMPLETE
PF	: TO COMPLETE
PSF	: TO COMPLETE
TSQ	: TO COMPLETE
P	: TO COMPLETE
PR	: TO COMPLETE
PT	: TO COMPLETE
tend	: TO COMPLETE

**Value**

data.frame with daily state variable

---

epirice.define.param *Define values of the parameters for the Epirice model*

---

**Description**

Define parameters values

**Usage**

```
epirice.define.param()
```

**Value**

matrix with parameter values (nominal, binf, bsup)

---

epirice.model                    *The Epirice model (Disease model for rice)*

---

### Description

**Model description.** Adapted from Savary et al.(2012)

### Usage

```
epirice.model(param, weather, sdate = 1, ldate = 120, H0 = 600)
```

### Arguments

param	a vector of parameters
weather	weather data.frame for one single year
sdate	date to begin simulation (day of year) (default 1)
ldate	date to end simulation (day of year) (default 120)
H0	initial number of plant's healthy sites(default 600)

### Value

data.frame with daily state variable

### See Also

[epirice.multi.simule](#)

---

epirice.multi.simule    *Wrapper function to run Epirice multiple times (for multiple sets of inputs)*

---

### Description

Wrapper function for epirice.model

### Usage

```
epirice.multi.simule(param, multi.simule, all = FALSE)
```

### Arguments

param	a vector of parameters
multi.simule	matrix of n row definition of input variable : site, year and date of transplantation.
all	if you want a matrix combining multi.simule and output (default = FALSE)

**Value**

matrix with AUDPC for each input vector

**See Also**

[epirice.model](#)

---

epirice.weather	<i>Read weather data for Epirice (southern Asia weather)</i>
-----------------	--

---

**Description**

Read weather data and format them for epirice.model

**Usage**

```
epirice.weather(working.year = NA, working.site = NA, weather = NA)
```

**Arguments**

working.year	year for the subset of weather data (default=NA : all the year)
working.site	site for the subset of weather data (default=NA : all the site)
weather	weather table

**Value**

data.frame with daily weather data for one or several site(s) and for one or several year(s)

---

evaluation.criteria	<i>Calcule multiple goodness-of-fit criteria</i>
---------------------	--

---

**Description**

This function is depreciated and will be remove from the package in future versions. Please use goodness.of.fit

**Usage**

```
evaluation.criteria(Ypred, Yobs, draw.plot = FALSE)
```

**Arguments**

Ypred	prediction values from the model
Yobs	observed values
draw.plot	draw evaluation plot

**Value**

data.frame with the different evaluation criteria

**Examples**

```
# observed and simulated values
obs<-c(78,110,92,75,110,108,113,155,150)
sim<-c(126,126,126,105,105,105,147,147,147)
evaluation.criteria(sim,obs,draw.plot=TRUE)
```

---

exponential.model      *The Exponential growth model of dynamic of population*

---

**Description**

Exponential growth model of dynamic of population

**Usage**

```
exponential.model(a, Y0, duration = 40, dt = 1)
```

**Arguments**

a                   : growth rate  
Y0                   : initial condition  
duration            : duration of simulation  
dt                   : time step for integration

**Value**

data.frame with Y for each time step

**See Also**

[verhulst.update](#) for the update function of the Verhulst model.

---

exponential.model.bis *The Exponential growth model of dynamic of population - another form*

---

**Description**

Exponential growth model of dynamic of population - another form

**Usage**

```
exponential.model.bis(a, Y0, duration = 40, dt = 1)
```

**Arguments**

a : growth rate  
Y0 : initial condition  
duration : duration of simulation  
dt : time step for integration

**Value**

data.frame with Y for each time step

**See Also**

[verhulst.update](#) for the update function of the Verhulst model.

---

exponential.model.ie *The Exponential growth model of dynamic of population - with improved Euler integration*

---

**Description**

Exponential growth model of dynamic of population - with improved Euler integration

**Usage**

```
exponential.model.ie(a, Y0, duration = 40, dt = 1)
```

**Arguments**

a : growth rate  
Y0 : initial condition  
duration : duration of simulation  
dt : time step for integration

**Value**

data.frame with Y for each time step

**See Also**

[verhulst.update](#) for the update function of the Verhulst model.

---

goodness.of.fit	<i>Calcule multiple goodness-of-fit criteria</i>
-----------------	--

---

**Description**

Calcule multiple goodness-of-fit criteria

**Usage**

```
goodness.of.fit(Yobs, Ypred, draw.plot = FALSE)
```

**Arguments**

Yobs	observed values
Ypred	prediction values from the model
draw.plot	draw evaluation plot

**Value**

data.frame with the different evaluation criteria

**Examples**

```
# observed and simulated values
obs<-c(78,110,92,75,110,108,113,155,150)
sim<-c(126,126,126,105,105,105,147,147,147)
goodness.of.fit(obs,sim,draw.plot=TRUE)
```

---

graph_epid	<i>Plot output of a Classical SEIR model for plant disease</i>
------------	--

---

**Description**

Plot the output of the Zadoks classical SEIR model for plant disease.

**Usage**

```
graph_epid(out, type1 = "s", all = TRUE, param = TRUE)
```

**Arguments**

out : output of the zadoks.original.model  
type1 : type of plot (default : s)  
all : if all=true (default), plot all the state variable  
param : if param (default), add the values of param on the plot

**Value**

plot

**See Also**

[zadoks.original.model](#)

---

graph_epid_s	<i>Plot output of a Classical SEIR model for plant disease</i>
--------------	--

---

**Description**

Plot the output of the Zadoks classical SEIR model for plant disease.

**Usage**

```
graph_epid_s(out, type1 = "s", all = TRUE, param = TRUE)
```

**Arguments**

out : output of the zadoks.original.model  
type1 : type of plot (default : s)  
all : if all=true (default), plot all the state variable  
param : if param (default), add the values of param on the plot

**Value**

plot

**See Also**[zakoks.original.model](#)

---

`lactation.calf.model` *The Lactation model*

---

**Description**

**Model description.** This model is a model of lactating mammary glands of cattle described by Heather et al. (1983). This model was then inspired more complex models based on these principles. This model simulates the dynamics of the production of cow's milk. the system is represented by 6 state variables: change in hormone levels (H), the production and loss of milk secreting cells (CS), and removing the secretion of milk (M), the average quantity of milk contained in the animal (Mmean), the amount of milk removed (RM) and yield (Y). The model has a time step  $dt = 0.1$  for regular consumption of milk by a calf. The model is defined by a few equations, with a total of fourteen parameters for the described process.

**Usage**

```
lactation.calf.model(  
  cu,  
  kdiv,  
  kd1,  
  kdh,  
  km,  
  ks1,  
  kr,  
  ks,  
  ksm,  
  mh,  
  mm,  
  p,  
  mum,  
  rc,  
  duration,  
  dt  
)
```

**Arguments**

cu : number of undifferentiated cells  
kdiv : cell division rate, Michaelis-Menten constant

kd1	: constant degradation of milk
kdh	: rate of decomposition of the hormone
km	: constant secretion of milk
ks1	: milk secretion rate, Michaelis-Menten constant
kr	: average milk constant
ks	: rate of degradation of the basal cells
ksm	: constant rate of degradation of milk secreting cells
mh	: parameter
mm	: storage Capacity milk the animal
p	: parameter
mum	: setting the maximum rate of cell division
rc	: parameter of milk m (t) function
duration	: duration of simulation
dt	: time step

**Value**

data.frame with CS, M, Mmoy, RM, day, week

**Examples**

```
lactation.calf.model2(lactation.define.param()["nominal",],300,0.1)
```

---

lactation.calf.model2 *The Lactation model for use with lactation.calf.simule*

---

**Description**

see lactation.calf.model for model description.

**Usage**

```
lactation.calf.model2(param, duration, dt)
```

**Arguments**

param	: a vector of parameters
duration	: duration of simulation
dt	: time step

**Value**

data.frame with CS, M, Mmoy, RM, day, week

**Examples**

```
sim=lactation.calf.model2(lactation.define.param()["nominal",],6+2*7, 0.1)
```

---

`lactation.calf.simule` *Wrapper function to run the Lactation model for multiple sets of parameter values*

---

### Description

Wrapper function to run the Lactation model for multiple sets of parameter values

### Usage

```
lactation.calf.simule(X, duration, dt)
```

### Arguments

`X` : parameter matrix  
`duration` : duration of simulation  
`dt` : time step

### Value

data.frame with : number of parameter vector (line number from X), week, CS, M, Mmoy, RM, day, week

---

`lactation.define.param`  
*Define values of the parameters for the Lactation model*

---

### Description

values from Heather et al. (1983) for different scenarios

### Usage

```
lactation.define.param(type = "calf")
```

### Arguments

`type` : for which model version ? "calf" or "machine"

### Value

matrix with parameter values (nominal, binf, bsup)

### Examples

```
lactation.define.param()
```

---

`lactation.machine.model`*The Lactation model with milking machine*

---

## Description

**Model description.** This model is a model of lactating mammary glands of cattle described by Heather et al. (1983). This model was then inspired more complex models based on these principles. This model simulates the dynamics of the production of cow's milk. the system is represented by 6 state variables: change in hormone levels (H), the production and loss of milk secreting cells (CS), and removing the secretion of milk (M), the average quantity of milk contained in the animal (Mmean), the amount of milk removed (RM) and yield (Y). The model has a time step  $dt = 0.001$  for milking machines. The model is defined by a few equations, with a total of twenty parameters for the described process.

## Usage

```
lactation.machine.model(  
  cu,  
  kdiv,  
  kd1,  
  kdh,  
  km,  
  ks1,  
  kr,  
  ks,  
  ksm,  
  mh,  
  mm,  
  p,  
  mum,  
  rma,  
  t1,  
  t2,  
  t3,  
  t4,  
  t5,  
  t6,  
  duration,  
  dt,  
  CSi,  
  Mi  
)
```

## Arguments

`cu` : number of undifferentiated cells

kdiv	: cell division rate, Michaelis-Menten constant
kd1	: constant degradation of milk
kdh	: rate of decomposition of the hormone
km	: constant secretion of milk
ks1	: milk secretion rate, Michaelis-Menten constant
kr	: average milk constant
ks	: rate of degradation of the basal cells
ksm	: constant rate of degradation of milk secreting cells
mh	: parameter
mm	: storage Capacity milk the animal
p	: parameter
mum	: setting the maximum rate of cell division
rma	: parameter of milk $m(t)$ function
t1	: parameter of milk $m(t)$ function
t2	: parameter of milk $m(t)$ function
t3	: parameter of milk $m(t)$ function
t4	: parameter of milk $m(t)$ function
t5	: parameter of milk $m(t)$ function
t6	: parameter of milk $m(t)$ function
duration	: duration of simulation
dt	: time step
CSi	: initial Number of secretory cells
Mi	: initial Quantity of milk in animal (kg)

**Value**

matrix with CS,M,Mmoy,RM

---

lactation.machine.model2

*The Lactation model for use with lactation.machine.simule*

---

**Description**

see lactation.calf.model for model description.

**Usage**

lactation.machine.model2(param, duration, dt, CSi, Mi)

**Arguments**

param	: a vector of parameters containing (cu,kdiv,kdl,kdh,km,ksl,kr,ks,ksm,mh,mm,p,mum,rma,t1,t2,t3,t4,t5,t6,t7,t8,t9,t10,t11,t12,t13,t14,t15,t16,t17,t18,t19,t20,t21,t22,t23,t24,t25,t26,t27,t28,t29,t30,t31,t32,t33,t34,t35,t36,t37,t38,t39,t40,t41,t42,t43,t44,t45,t46,t47,t48,t49,t50,t51,t52,t53,t54,t55,t56,t57,t58,t59,t60,t61,t62,t63,t64,t65,t66,t67,t68,t69,t70,t71,t72,t73,t74,t75,t76,t77,t78,t79,t80,t81,t82,t83,t84,t85,t86,t87,t88,t89,t90,t91,t92,t93,t94,t95,t96,t97,t98,t99,t100,t101,t102,t103,t104,t105,t106,t107,t108,t109,t110,t111,t112,t113,t114,t115,t116,t117,t118,t119,t120,t121,t122,t123,t124,t125,t126,t127,t128,t129,t130,t131,t132,t133,t134,t135,t136,t137,t138,t139,t140,t141,t142,t143,t144,t145,t146,t147,t148,t149,t150,t151,t152,t153,t154,t155,t156,t157,t158,t159,t160,t161,t162,t163,t164,t165,t166,t167,t168,t169,t170,t171,t172,t173,t174,t175,t176,t177,t178,t179,t180,t181,t182,t183,t184,t185,t186,t187,t188,t189,t190,t191,t192,t193,t194,t195,t196,t197,t198,t199,t200,t201,t202,t203,t204,t205,t206,t207,t208,t209,t210,t211,t212,t213,t214,t215,t216,t217,t218,t219,t220,t221,t222,t223,t224,t225,t226,t227,t228,t229,t230,t231,t232,t233,t234,t235,t236,t237,t238,t239,t240,t241,t242,t243,t244,t245,t246,t247,t248,t249,t250,t251,t252,t253,t254,t255,t256,t257,t258,t259,t260,t261,t262,t263,t264,t265,t266,t267,t268,t269,t270,t271,t272,t273,t274,t275,t276,t277,t278,t279,t280,t281,t282,t283,t284,t285,t286,t287,t288,t289,t290,t291,t292,t293,t294,t295,t296,t297,t298,t299,t300,t301,t302,t303,t304,t305,t306,t307,t308,t309,t310,t311,t312,t313,t314,t315,t316,t317,t318,t319,t320,t321,t322,t323,t324,t325,t326,t327,t328,t329,t330,t331,t332,t333,t334,t335,t336,t337,t338,t339,t340,t341,t342,t343,t344,t345,t346,t347,t348,t349,t350,t351,t352,t353,t354,t355,t356,t357,t358,t359,t360,t361,t362,t363,t364,t365,t366,t367,t368,t369,t370,t371,t372,t373,t374,t375,t376,t377,t378,t379,t380,t381,t382,t383,t384,t385,t386,t387,t388,t389,t390,t391,t392,t393,t394,t395,t396,t397,t398,t399,t400,t401,t402,t403,t404,t405,t406,t407,t408,t409,t410,t411,t412,t413,t414,t415,t416,t417,t418,t419,t420,t421,t422,t423,t424,t425,t426,t427,t428,t429,t430,t431,t432,t433,t434,t435,t436,t437,t438,t439,t440,t441,t442,t443,t444,t445,t446,t447,t448,t449,t450,t451,t452,t453,t454,t455,t456,t457,t458,t459,t460,t461,t462,t463,t464,t465,t466,t467,t468,t469,t470,t471,t472,t473,t474,t475,t476,t477,t478,t479,t480,t481,t482,t483,t484,t485,t486,t487,t488,t489,t490,t491,t492,t493,t494,t495,t496,t497,t498,t499,t500,t501,t502,t503,t504,t505,t506,t507,t508,t509,t510,t511,t512,t513,t514,t515,t516,t517,t518,t519,t520,t521,t522,t523,t524,t525,t526,t527,t528,t529,t530,t531,t532,t533,t534,t535,t536,t537,t538,t539,t540,t541,t542,t543,t544,t545,t546,t547,t548,t549,t550,t551,t552,t553,t554,t555,t556,t557,t558,t559,t560,t561,t562,t563,t564,t565,t566,t567,t568,t569,t570,t571,t572,t573,t574,t575,t576,t577,t578,t579,t580,t581,t582,t583,t584,t585,t586,t587,t588,t589,t590,t591,t592,t593,t594,t595,t596,t597,t598,t599,t600,t601,t602,t603,t604,t605,t606,t607,t608,t609,t610,t611,t612,t613,t614,t615,t616,t617,t618,t619,t620,t621,t622,t623,t624,t625,t626,t627,t628,t629,t630,t631,t632,t633,t634,t635,t636,t637,t638,t639,t640,t641,t642,t643,t644,t645,t646,t647,t648,t649,t650,t651,t652,t653,t654,t655,t656,t657,t658,t659,t660,t661,t662,t663,t664,t665,t666,t667,t668,t669,t670,t671,t672,t673,t674,t675,t676,t677,t678,t679,t680,t681,t682,t683,t684,t685,t686,t687,t688,t689,t690,t691,t692,t693,t694,t695,t696,t697,t698,t699,t700,t701,t702,t703,t704,t705,t706,t707,t708,t709,t710,t711,t712,t713,t714,t715,t716,t717,t718,t719,t720,t721,t722,t723,t724,t725,t726,t727,t728,t729,t730,t731,t732,t733,t734,t735,t736,t737,t738,t739,t740,t741,t742,t743,t744,t745,t746,t747,t748,t749,t750,t751,t752,t753,t754,t755,t756,t757,t758,t759,t760,t761,t762,t763,t764,t765,t766,t767,t768,t769,t770,t771,t772,t773,t774,t775,t776,t777,t778,t779,t780,t781,t782,t783,t784,t785,t786,t787,t788,t789,t790,t791,t792,t793,t794,t795,t796,t797,t798,t799,t800,t801,t802,t803,t804,t805,t806,t807,t808,t809,t810,t811,t812,t813,t814,t815,t816,t817,t818,t819,t820,t821,t822,t823,t824,t825,t826,t827,t828,t829,t830,t831,t832,t833,t834,t835,t836,t837,t838,t839,t840,t841,t842,t843,t844,t845,t846,t847,t848,t849,t850,t851,t852,t853,t854,t855,t856,t857,t858,t859,t860,t861,t862,t863,t864,t865,t866,t867,t868,t869,t870,t871,t872,t873,t874,t875,t876,t877,t878,t879,t880,t881,t882,t883,t884,t885,t886,t887,t888,t889,t890,t891,t892,t893,t894,t895,t896,t897,t898,t899,t900,t901,t902,t903,t904,t905,t906,t907,t908,t909,t910,t911,t912,t913,t914,t915,t916,t917,t918,t919,t920,t921,t922,t923,t924,t925,t926,t927,t928,t929,t930,t931,t932,t933,t934,t935,t936,t937,t938,t939,t940,t941,t942,t943,t944,t945,t946,t947,t948,t949,t950,t951,t952,t953,t954,t955,t956,t957,t958,t959,t960,t961,t962,t963,t964,t965,t966,t967,t968,t969,t970,t971,t972,t973,t974,t975,t976,t977,t978,t979,t980,t981,t982,t983,t984,t985,t986,t987,t988,t989,t990,t991,t992,t993,t994,t995,t996,t997,t998,t999,1000)
duration	: duration of simulation
dt	: time step
CSi	: initial Number of secretory cells
Mi	: initial Quantity of milk in animal (kg)

**Value**

data.frame with CS, M, Mmoy, RM, day, week

---

magarey.define.param *Define values of the parameters for the Magarey model*

---

**Description**

Define values of the parameters for the Magarey model

**Usage**

```
magarey.define.param(species = "unkown")
```

**Arguments**

species : name of a species. By default, value for an "unkown" species are given. Other possibility are "G.citricarpa" or "Kaki.fungus"

**Value**

matrix with parameter values (nominal, binf, bsup)

**Examples**

```
magarey.define.param(species="G.citricarpa")
magarey.define.param(species="Kaki.fungus")
```

---

 magarey.model

*The Magarey model*


---

**Description**

Generic model of infection for foliar diseases caused by fungi (from Magarey et al.,2005).

**Usage**

```
magarey.model(T, Tmin, Topt, Tmax, Wmin, Wmax)
```

**Arguments**

T : input variable. Either a scalar or a vector (for a weather series).  
 Tmin : parameter of minimal temperature for infection (degC)  
 Topt : parameter of optimal temperature for infection (degC)  
 Tmax : parameter of maximal temperature for infection (degC)  
 Wmin : parameter of minimal wetness duration for infection (hour)  
 Wmax : parameter of maximal wetness duration for infection (hour)

**Value**

Wetness duration (W, hour). Either a scalar or a vector depending on T.

**Examples**

```
plot(1:35, magarey.model (1:35,7, 18, 30, 10, 42), type="l", xlab="T", ylab="W")
```

---

 magarey.model2

*The Magarey model, taking a vector of parameters as argument*


---

**Description**

Generic model of infection for foliar diseases caused by fungi (from Magarey et al.,2005).

**Usage**

```
magarey.model2(T, param)
```

**Arguments**

T : input variable. Either a scalar or a vector (for a weather series).  
 param : parameters

**Value**

W : Wetness duration (hour). Either a scalar or a vector depending on T.

---

magarey.simule	<i>Wrapper function to run the Magarey model multiple times (for multiple sets of inputs)</i>
----------------	---

---

**Description**

Wrapper function to run the Magarey model multiple times (for multiple sets of inputs)

Example `magarey.simule(magarey.define.param(),15)`

**Usage**

```
magarey.simule(X, T, all = FALSE)
```

**Arguments**

X : parameter matrix  
T : input variable, temperature  
all : if you want a matrix combining X and output

**Value**

a table with wetness duration (W) for each parameter vector

---

maize.data_EuropeEU	<i>Maize Biomass and Leaf Area Data</i>
---------------------	---

---

**Description**

Observation data for several sites and years in Europe. This data are fake observed data, derived from simulations with an error model.

**Usage**

```
data(maize.data_EuropeEU)
```

**Format**

A RangedData instance, 1 row per measurement.

**Source**

NA

maize.data\_MetaModelling  
*dataset of simulation for maize final biomass*

---

### Description

Simulation data for several site and year (site-year) in Europe. This data are from run of the original maize crop model with the R function `maize.model` for the 30 French sites and 17 years included in the dataset `weather_FranceWest` of the package `ZeBook`. Our training dataset includes 510 (30 sites \* 17 years) simulated biomass values and the 510 corresponding series of input values. The input values are the three average temperatures T1, T2, T3 and the three average radiations RAD1, RAD2, RAD3 computed on 3 periods of the growing season. Period 1 : from day 1 to day 50 (day of the year), period 2: from day 51 to day 100, period 3 : from day 101 to day 150.

### Usage

```
data(maize.data_EuropeEU)
```

### Format

a `RangedData` instance, 1 row per simulation. Site, Year, T1, T2, T3, RAD1, RAD2, RAD3, B

### Source

NA

### See Also

[maize.model](#), [weather\\_FranceWest](#)

---

maize.define.param     *Define values of the parameters for the Maize model*

---

### Description

Define parameters values

### Usage

```
maize.define.param()
```

### Value

matrix with parameter values (nominal, binf, bsup)

---

maize.model	<i>The basic Maize model.</i>
-------------	-------------------------------

---

### Description

**Model description.** This model is a dynamic model of crop growth for Maize cultivated in potential conditions. The crop growth is represented by three state variables, leaf area per unit ground area (leaf area index, LAI), total biomass (B) and cumulative thermal time since plant emergence (TT). It is based on key concepts included in most crop models, at least for the "potential production" part. In fact, this model does not take into account any effects of soil water, nutrients, pests, or diseases,...

### Usage

```
maize.model(Tbase, RUE, K, alpha, LAImax, TTM, TTL, weather, sdate, ldate)
```

### Arguments

Tbase	parameter the baseline temperature for growth (degreeCelsius)
RUE	parameter radiation use efficiency (?)
K	parameter extinction coefficient (relation between leaf area index and intercepted radiation) (-)
alpha	parameter the relative rate of leaf area index increase for small values of leaf area index (?)
LAImax	parameter maximum leaf area index (-)
TTM	parameter temperature sum for crop maturity (degreeC.day)
TTL	parameter temperature sum at the end of leaf area increase (degreeC.day)
weather	weather data.frame for one single year
sdate	sowing date
ldate	last date

### Details

The tree state variables are dynamic variables depending on days after emergence:  $TT(day)$ ,  $B(day)$ , and  $LAI(day)$ . The model has a time step  $dt$  of one day.

The model is defined by a few equations, with a total of seven parameters for the described process.

$$(1) TT(day + 1) = TT(day) + dTT(day)$$

$$(2) B(day + 1) = B(day) + dB(day)$$

$$(3) LAI(day + 1) = LAI(day) + dLAI(day)$$

$$(4) dTT(day) = \max\left(\frac{TMIN(day) + TMAX(day)}{2} - Tbase; 0\right)$$

$$(5) dB(day) = RUE * (1 - e^{-K * LAI(day) * I(day)}), \text{ if } TT(day) \leq TTM$$

$$dB(day) = 0, \text{ if } TT(day) > TTM$$

$$(6) dLAI(day) = alpha * dTT(day) * LAI(day) * \max(LAImax - LAI(day); 0), \text{ if } TT(day) \leq TTL$$

$$dLAI(day) = 0, \text{ if } TT(day) > TTL$$

**Value**

data.frame with daily TT, LAI,B

**See Also**

[maize.model2](#), [maize.define.param](#), [maize.simule](#), [maize.multisy](#), [maize.simule240](#), [maize.simule\\_multisy240](#)

**Examples**

```
weather = maize.weather(working.year=2010, working.site=30,weather_all=weather_EuropeEU)
maize.model(Tbase=7, RUE=1.85, K=0.7, alpha=0.00243, LAImax=7, TTM=1200, TTL=700,
            weather, sdate=100, ldate=250)
```

---

maize.model2

*The basic Maize model for use with maize.simule*

---

**Description**

Wrapper pour maize.model

**Usage**

```
maize.model2(param, weather, sdate, ldate)
```

**Arguments**

param : a vector of parameters  
weather : weather data.frame for one single year  
sdate : sowing date  
ldate : last date

**Value**

data.frame with daily TT, LAI,B

**Examples**

```
weather = maize.weather(working.year=2010, working.site=30,weather_all=weather_EuropeEU)
maize.model2(maize.define.param()["nominal",], weather, sdate=100, ldate=250)
```

---

maize.muchow.graph      *Plot dynamic output of Muchow Maize model.*

---

**Description**

Plot 6 graphs of main output variables of the Muchow Maize model.

**Usage**

```
maize.muchow.graph(res)
```

**Arguments**

res                    list of result from maize.muchow.model

**See Also**

[mm.A.fct](#), [mm.LN.fct](#), [mm.FAS.fct](#), [maize.multisy](#), [mm.HI.fct](#), [maize.muchow.model](#)

**Examples**

```
# not run in package test
# res = maize.muchow.model(weather=maize.weather(working.year=2010, working.site=1))
# maize.muchow.graph(res)
```

---

maize.muchow.model      *Maize model, with harvest index and yield from Muchow et al. (1990)*

---

**Description**

Maize model, with harvest index and yield. from Muchow RC, Sinclair TR, and Bennett JM (1990). Temperature and Solar Radiation Effects on Potential Maize Yield across Locations AGRONOMY JOURNAL, VOL. 82, MARCH-APRIL 1990

**Usage**

```
maize.muchow.model(  
  Tbase1 = 8,  
  TTE = 87,  
  TTS = 67,  
  Tbase2 = 0,  
  TTRUE = 500,  
  TTM = 1150,  
  TLN = 20,  
  AM = 596,  
  RUE1 = 1.6,
```

```

RUE2 = 1.2,
K = 0.4,
HImax = 0.5,
Population = 7,
sdate = 100,
ldate = 365,
weather
)

```

### Arguments

Tbase1	base temperature before silking(degC)
TTE	Thermal units from sowing to emergence/leaf growth (degC.day)
TTS	Thermal units from end of leaf growth to silking (degC.day)
Tbase2	base temperature after silking (degC)
TTRUE	Thermal units from silking for RUE change (degC.day)
TTM	Thermal units from silking to physiological maturity (degC.day)
TLN	total number of leaves initiated (-)
AM	area of the largest leaf (cm2)
RUE1	radiation use efficiency (g.MJ-1) from crop emergence until 500 thermal units (base 0 "C) after silking
RUE2	radiation use efficiency (g.MJ-1) from 500 thermal units (base 0 "C) after silking
K	radiation extinction coefficient (-)
HImax	maximum harvest index - genetic potential (-)
Population	number of plant per square meter (-)
sdate	sowing date (day)
ldate	end of simulation (day)
weather	daily weather dataframe

### Value

data.frame with TT1, TT2, STADE, LN, LAI, B, HI, YIELD

### See Also

[mm.A.fct](#), [mm.LN.fct](#), [mm.FAS.fct](#), [maize.multisy](#), [mm.HI.fct](#), [maize.muchow.graph](#)

### Examples

```

# not run in package test
# res = maize.muchow.model(weather=maize.weather(working.year=2010, working.site=1))
#res$FinalYield

```

---

maize.multisy	<i>Wrapping function to run maize model on several site-years</i>
---------------	---

---

**Description**

Wrapping function to run maize model on several site-years

**Usage**

```
maize.multisy(param, list_site_year, sdate, ldate, weather_all = NA)
```

**Arguments**

param : a vector of parameters  
list\_site\_year : vector of site-year  
sdate : sowing date  
ldate : last date  
weather\_all : weather data.frame for corresponding site-years

**Value**

a data.frame with simulation for all site-years, with the first column sy indicating the site-years

---

maize.multisy240	<i>Wrapper function to run Maize model for multiple sets of input variables (site-year) and give Biomass at day240.</i>
------------------	---

---

**Description**

Wrapper function to run Maize model for multiple sets of input variables (site-year) and give Biomass at day240.

**Usage**

```
maize.multisy240(param, liste_sy, sdate, ldate, weather_all = NA)
```

**Arguments**

param : a vector of parameters  
liste\_sy : vector of site-year  
sdate : sowing date  
ldate : last date  
weather\_all : weather data table used

**Value**

mean biomass at day=240

**Examples**

```
maize.multisy240(maize.define.param()["nominal"],c("18-2006","64-2004")
, sdate=100, ldate=250, weather_all=weather_EuropeEU)
```

---

maize.RUEtemp

*Calculate effect of temperature on RUE for Maize*

---

**Description**

Function to compute effect of temperature on RUE

**Usage**

```
maize.RUEtemp(T, RUE_max, T0, T1, T2, T3)
```

**Arguments**

T : temperature  
RUE\_max : maximum value for RUE  
T0 : temperature parameter  
T1 : temperature parameter  
T2 : temperature parameter  
T3 : temperature parameter

**Value**

RUE value

---

maize.simule

*Wrapper function to run Maize model for multiple sets of parameter values*

---

**Description**

wrapper for maize.model2

**Usage**

```
maize.simule(X, weather, sdate, ldate, all = FALSE)
```

**Arguments**

X : matrix of n row vectors of 7 parameters  
 weather : weather data.frame for one single year  
 sdate : sowing date  
 ldate : last date  
 all : if you want a matrix combining X and output (default = FALSE)

**Value**

matrix with maximum biomass for each parameter vector

---

maize.simule240	<i>Wrapper function to run Maize model multiple times for multiple sets of parameter values and give Biomass at day240</i>
-----------------	--

---

**Description**

Wrapper function for multiple simulation with Maize model

**Usage**

```
maize.simule240(X, weather, sdate, ldate, all = FALSE)
```

**Arguments**

X : matrix of n row vectors of 7 parameters  
 weather : weather data.frame for one single year  
 sdate : sowing date  
 ldate : last date  
 all : if you want a matrix combining X and output (default = FALSE)

**Value**

a matrix of biomass at day=240 for all combinations of parameters of X

**Examples**

```

sy="18-2006"
weather = maize.weather(working.year=strsplit(sy,"-")[1][2],
  working.site=strsplit(sy,"-")[1][1],weather_all=weather_EuropeEU)
maize.simule240(maize.define.param(),weather, sdate=100, ldate=250, all=FALSE)

```

---

```
maize.simule_multisy240
```

*Wrapper function to run Maize model for multiple sets of parameter values (virtual design) and multiple sets of input variables (site-year) and give Biomass at day240*

---

### Description

Wrapper function to run Maize model for multiple sets of input variables (site-year) and give Biomass at day240.

### Usage

```
maize.simule_multisy240(  
  X,  
  liste_sy,  
  sdate,  
  ldate,  
  weather_all = NA,  
  all = FALSE  
)
```

### Arguments

X	matrix of n row vectors of 7 parameters
liste_sy	vector of site-year
sdate	sowing date
ldate	last date
weather_all	Weather data base
all	if you want a matrix combining X and output (default = FALSE)

### Value

a matrix of mean biomass at day=240 for all combinations of parameters of X

### Examples

```
maize.simule_multisy240(maize.define.param(),c("18-2006", "64-2004"),  
  sdate=100, ldate=250,weather_all=weather_EuropeEU,all=FALSE)
```

---

maize.weather	<i>Read weather data for the Maize model</i>
---------------	--

---

**Description**

Function to read weather data and format them for maize.model

**Usage**

```
maize.weather(working.year = NA, working.site = NA, weather_all = NA)
```

**Arguments**

working.year	year for the subset of weather data (default=NA : all the year)
working.site	site for the subset of weather data (default=NA : all the site)
weather_all	weather data base (default=weather_FranceWest)

**Value**

data.frame with daily weather data for one or several site(s) and for one or several year(s)

---

maize_cir.model	<i>The Maize model with additional state variable CumInt</i>
-----------------	--

---

**Description**

Variant of the maize model

**Usage**

```
maize_cir.model(Tbase, RUE, K, alpha, LAImax, TTM, TTL, weather, sdate, ldate)
```

**Arguments**

Tbase	parameter the baseline temperature for growth (degreeCelsius)
RUE	parameter radiation use efficiency (?)
K	parameter extinction coefficient (relation between leaf area index and intercepted radiation) (-)
alpha	parameter the relative rate of leaf area index increase for small values of leaf area index (?)
LAImax	parameter maximum leaf area index (-)
TTM	parameter temperature sum for crop maturity (degreeC.day)
TTL	parameter temperature sum at the end of leaf area increase (degreeC.day)
weather	weather data.frame for one single year
sdate	sowing date
ldate	last date

**Value**

data.frame with daily TT, LAI,B

---

maize\_cir\_rue.model    *The Maize model with temperature dependent RUE and CumInt*

---

**Description**

Variant of the maize.model

**Usage**

```
maize_cir_rue.model(
  Tbase,
  RUE_max,
  K,
  alpha,
  LAImax,
  TTM,
  TTL,
  weather,
  sdate,
  ldate
)
```

**Arguments**

Tbase	parameter the baseline temperature for growth (degreeCelsius)
RUE_max	parameter maximum radiation use efficiency (?)
K	parameter extinction coefficient (relation between leaf area index and intercepted radiation) (-)
alpha	parameter the relative rate of leaf area index increase for small values of leaf area index (?)
LAImax	parameter maximum leaf area index (-)
TTM	parameter temperature sum for crop maturity (degreeC.day)
TTL	parameter temperature sum at the end of leaf area increase (degreeC.day)
weather	weather data.frame for one single year
sdate	sowing date
ldate	last date

**Value**

data.frame with daily TT, LAI,B

---

 maize\_cir\_rue\_ear.model

*The Maize model with temperature dependent RUE, CumInt and ear growth*

---

## Description

Variant of the maize.model

## Usage

```
maize_cir_rue_ear.model(  
  Tbase,  
  RUE_max,  
  K,  
  alpha,  
  LAImax,  
  TTM,  
  TTL,  
  weather,  
  sdate,  
  ldate  
)
```

## Arguments

Tbase	parameter the baseline temperature for growth (degreeCelsius)
RUE_max	parameter maximum radiation use efficiency (?)
K	parameter extinction coefficient (relation between leaf area index and intercepted radiation) (-)
alpha	parameter the relative rate of leaf area index increase for small values of leaf area index (?)
LAImax	parameter maximum leaf area index (-)
TTM	parameter temperature sum for crop maturity (degreeC.day)
TTL	parameter temperature sum at the end of leaf area increase (degreeC.day)
weather	weather data.frame for one single year
sdate	sowing date
ldate	last date

## Value

data.frame with daily TT, LAI,B

---

 mm.A.fct

*Expanded leaf area function for Muchow et al. (1990) Maize model*


---

**Description**

Compute fully expanded area by leaf number (A, cm<sup>2</sup>)

**Usage**

```
mm.A.fct(LN, AM, LNM, a1 = -0.0344, a2 = 0.000731)
```

**Arguments**

LN	Leaf number
AM	area of the largest leaf (cm <sup>2</sup> )
LNM	leaf number having the largest area (-)
a1	coefficient of the statistical relation (default : -0.0344)
a2	coefficient of the statistical relation (default : 0.000731)

**Value**

vector of Expanded leaf area

**See Also**

[maize.muchow.model](#), [mm.LN.fct](#), [mm.FAS.fct](#), [maize.multisy](#), [mm.HI.fct](#), [maize.muchow.graph](#)

**Examples**

```
barplot(mm.A.fct(LN=1:20, AM=750, LNM=12), names.arg=1:20,
        horiz=TRUE, xlab="leaf area (cm2)", ylab="leaf number")
```

---

 mm.FAS.fct

*Senescence function for Muchow et al. (1990) Maize model*


---

**Description**

Senesced fraction of total leaf area (FAS) increase with thermal units (TU) from emergence

**Usage**

```
mm.FAS.fct(TT, TTE, c1 = 0.00161, c2 = 0.00328)
```

**Arguments**

TT	Thermal time (degC.day)
TTE	Thermal units from sowing to emergence/leaf growth (degC.day)
c1	coefficient of the statistical relation (default : 0.00161)
c2	coefficient of the statistical relation (default : 0.00328)

**Value**

Senesced fraction of total leaf area

**See Also**

[maize.muchow.model](#), [mm.A.fct](#), [mm.LN.fct](#), [maize.multisy](#), [mm.HI.fct](#), [maize.muchow.graph](#)

**Examples**

```
plot(1:2500, mm.FAS.fct(1:2500, TTE=87))
```

---

mm.HI.fct

*Harvest index function for Muchow et al. (1990) Maize model*

---

**Description**

Compute the harvest index.

**Usage**

```
mm.HI.fct(day, daysilking, HImax, d1 = 0.015, d2 = 3)
```

**Arguments**

day	day of the year
daysilking	day of the year for silking (day)
HImax	maximum harvest index - genetic potential (-)
d1	coefficient of the statistical relation (day-1, default : 0.015)
d2	coefficient of the statistical relation (day, default : 3)

**Value**

Harvest index

**See Also**

[maize.muchow.model](#), [mm.A.fct](#), [mm.LN.fct](#), [maize.multisy](#), [mm.FAS.fct](#), [maize.muchow.graph](#)

**Examples**

```
plot(1:350, mm.HI.fct(1:350, 200, 0.75), type="l")
```

---

 mm.LN.fct

*Leaf number function for Muchow et al. (1990) Maize model*


---

**Description**

Leaf number as a function of thermal time

**Usage**

```
mm.LN.fct(TT1, TTE, b1 = 2.5, b2 = 0.00225, TLN = 20)
```

**Arguments**

TT1	Thermal time from sowing (degC.day)
TTE	Thermal units from sowing to emergence/leaf growth (degC.day)
b1	coefficient of the statistical relation (default : 2.5)
b2	coefficient of the statistical relation (default : 0.00225)
TLN	total number of leaves initiated (-)

**Value**

Leaf number

**See Also**

[maize.muchow.model](#), [mm.A.fct](#), [mm.FAS.fct](#), [maize.multisy](#), [mm.HI.fct](#), [maize.muchow.graph](#)

**Examples**

```
plot(1:1000, mm.LN.fct(1:1000, TTE=87))
```

---

 param.rtriangle

*Generate a random plan as a data frame. Columns are parameters. Values have triangle distribution*


---

**Description**

according to nominal, minimal and maximal values defined in a model.factors matrix

**Usage**

```
param.rtriangle(model.factors, N)
```

**Arguments**

model.factors : matrix defining nominal, minimal (binf), maximal values (bsup) for a set of p parameters  
 N : size of sample

**Value**

parameter matrix of dim = (N, p)

---

param.runif                      *Generate a random plan as a data frame. Columns are parameters. Values have uniform distribution*

---

**Description**

according to minimal and maximal values defined in a model.factors matrix

**Usage**

```
param.runif(model.factors, N)
```

**Arguments**

model.factors : matrix defining minimal (binf) and maximal values (bsup) for a set of p parameters  
 N : size of sample

**Value**

parameter matrix of dim = (N, p)

---

population.age.matrix.model  
*The PopulationAge model (Population Dynamics with Age Classes) - matrix form*

---

**Description**

Population Dynamics Model with Age Classes for an insect Exactly the same model as population.age.model, but written as a matrix computation. It's possible for this model. It's really more efficient and reduce computer time by 6! 7 states variables E : egg stage. homogenous population (density) (number per ha) L1 : larvae1 stage. homogenous population (density) (number per ha) L2 : larvae2 stage. homogenous population (density) (number per ha) L3 : larvae3 stage. homogenous population (density) (number per ha) L4 : larvae4 stage. homogenous population (density) (number per ha) P : pupae stage. homogenous population (density) (number per ha) A : adult stage. homogenous population (density) (number per ha)

**Usage**

```

population.age.matrix.model(
  rb = 3.5,
  mE = 0.017,
  rE = 0.172,
  m1 = 0.06,
  r12 = 0.217,
  m2 = 0.032,
  r23 = 0.313,
  m3 = 0.022,
  r34 = 0.222,
  m4 = 0.02,
  r4P = 0.135,
  mP = 0.02,
  rPA = 0.099,
  mA = 0.027,
  iA = 0,
  duration = 100,
  dt = 1
)

```

**Arguments**

rb	: eggs laid per adult per unit area (day-1)
mE	: relative mortality rate of egg (day-1)
rE	: eggs hatch (day-1)
m1	: relative mortality rate of larvae L1 (day-1)
r12	: relative rate L1->L2 (day-1)
m2	: relative mortality rate of larvae L2 (day-1)
r23	: relative rate L2->L3 (day-1)
m3	: relative mortality rate of larvae L3 (day-1)
r34	: relative rate L3->L4 (day-1)
m4	: relative mortality rate of larvae L4 (day-1)
r4P	: relative rate L4->P (day-1)
mP	: relative mortality rate of pupae (day-1)
rPA	: relative rate P->A (day-1)
mA	: relative mortality rate of adult L1 (day-1)
iA	: input rate of adult (unit.day-1)
duration	: simulation duration
dt	: time step for integration

**Value**

data.frame with values for state variables for each time step.

---

population.age.model    *The PopulationAge model (Population Dynamics with Age Classes)*

---

## Description

Population Dynamics Model with Age Classes for an insect

## Usage

```
population.age.model(
  rb = 3.5,
  mE = 0.017,
  rE = 0.172,
  m1 = 0.06,
  r12 = 0.217,
  m2 = 0.032,
  r23 = 0.313,
  m3 = 0.022,
  r34 = 0.222,
  m4 = 0.02,
  r4P = 0.135,
  mP = 0.02,
  rPA = 0.099,
  mA = 0.027,
  iA = 0,
  duration = 100,
  dt = 1
)
```

## Arguments

rb	eggs laid per adult per unit area (day-1)
mE	relative mortality rate of egg (day-1)
rE	eggs hatch (day-1)
m1	relative mortality rate of larvae L1 (day-1)
r12	relative rate L1->L2 (day-1)
m2	relative mortality rate of larvae L2 (day-1)
r23	relative rate L2->L3 (day-1)
m3	relative mortality rate of larvae L3 (day-1)
r34	relative rate L3->L4 (day-1)
m4	relative mortality rate of larvae L4 (day-1)
r4P	relative rate L4->P (day-1)
mP	relative mortality rate of pupae (day-1)

rPA	relative rate P->A (day-1)
mA	relative mortality rate of adult L1 (day-1)
iA	input rate of adult (unit.day-1)
duration	simulation duration
dt	time step for integration

**Value**

data.frame with values for state variables for each time step.

---

population.age.model.ode

*The PopulationAge model (Population Dynamics with Age Classes) -  
ode form*

---

**Description**

Population Dynamics Model with Age Classes for an insect Exactly the same model as population.age.model, but written as an ordinary differential equation system (ode) with deSolve package. 7 states variables E egg stage. homogenous population (density) (number per ha) L1 larvae1 stage. homogenous population (density) (number per ha) L2 larvae2 stage. homogenous population (density) (number per ha) L3 larvae3 stage. homogenous population (density) (number per ha) L4 larvae4 stage. homogenous population (density) (number per ha) P pupae stage. homogenous population (density) (number per ha) A adult stage. homogenous population (density) (number per ha)

**Usage**

```
population.age.model.ode(
  rb = 3.5,
  mE = 0.017,
  rE = 0.172,
  m1 = 0.06,
  r12 = 0.217,
  m2 = 0.032,
  r23 = 0.313,
  m3 = 0.022,
  r34 = 0.222,
  m4 = 0.02,
  r4P = 0.135,
  mP = 0.02,
  rPA = 0.099,
  mA = 0.027,
  iA = 0,
  duration = 100,
  dt = 1,
  method = "euler"
)
```

**Arguments**

rb	eggs laid per adult per unit area (day-1)
mE	relative mortality rate of egg (day-1)
rE	eggs hatch (day-1)
m1	relative mortality rate of larvae L1 (day-1)
r12	relative rate L1->L2 (day-1)
m2	relative mortality rate of larvae L2 (day-1)
r23	relative rate L2->L3 (day-1)
m3	relative mortality rate of larvae L3 (day-1)
r34	relative rate L3->L4 (day-1)
m4	relative mortality rate of larvae L4 (day-1)
r4P	relative rate L4->P (day-1)
mP	relative mortality rate of pupae (day-1)
rPA	relative rate P->A (day-1)
mA	relative mortality rate of adult L1 (day-1)
iA	input rate of adult (unit.day-1)
duration	simulation duration
dt	time step for integration
method	integration method (euler, rk4,...)

**Value**

data.frame with values for state variables for each time step.

---

predator.prey.model    *The Predator-Prey model (Predator-Prey Lotka-Volterra with logistic equation for prey)*

---

**Description**

Predator-Prey Lotka-Volterra model (with logistic prey)

**Usage**

```
predator.prey.model(
  grH = 1,
  kH = 10,
  mrH = 0.2,
  eff = 0.5,
  mrA = 0.2,
  H0 = 1,
```

```

A0 = 2,
duration = 200,
dt = 1,
method = "euler"
)

```

### Arguments

grH : relative rate of prey population growth  
 kH : environment carrying capacity for prey (number per ha)  
 mrH : maximum predation rate (number per predator and per prey per day)  
 eff : efficiency, growth of predator population depending on predation (-)  
 mrA : mortality of predator (-)  
 H0 : size of population of prey, at time 0  
 A0 : size of population of predator, at time 0  
 duration : simulation duration  
 dt : time step for integration  
 method : integration method

### Value

data.frame with daily H and A

---

q.arg.fast.runif      *Build the q.arg argument for the FAST function (sensitivity analysis)*

---

### Description

according to minimal and maximal values defined in a model.factors matrix

### Usage

```
q.arg.fast.runif(model.factors)
```

### Arguments

model.factors : matrix defining minimal (binf) and maximal values (bsup) for a set of p parameters

### Value

a list of list

---

seedweight.data	<i>Wheat grain weight measurements after anthesis</i>
-----------------	---

---

**Description**

Darroch and Baker (1990) studied grain filling in three spring wheat genotypes. The data are seed weights of the spring wheat cultivar Neepawa in three different years 1986, 1987, 1988. These data were numerised from figure of the article, so they present slight difference with original data.

**Usage**

```
seedweight.data
```

**Format**

a RangedData instance, 1 row per measurement. DD = Degree Days after anthesis; seedweight = Wheat grain weight (mg)

**Source**

Darroch, B A, and R J Baker. 1990. "Grain filling in three spring wheat genotypes: statistical analysis." *Crop Science* 30(3):525-529.

---

seedweight.model	<i>The SeedWeight model</i>
------------------	-----------------------------

---

**Description**

The SeedWeight model is a logistic model of grain weight over time in wheat. The model was proposed by Darroch & Baker (1990) in a study of grain filling in three spring wheat genotypes. This model has a single input variable, degree days after anthesis noted DD, and three parameters, noted W, B and C. Parameters are estimated from observations.

**Usage**

```
seedweight.model(DD, W, B, C)
```

**Arguments**

DD	: degree days after anthesis
W	: parameter of the model
B	: parameter of the model
C	: parameter of the model

**Value**

Seed Weight for each TT

**Examples**

```
plot(1:500, seedweight.model(1:500, W=30, B=4, C=0.020), type="l",
     xlab="degree days after anthesis", ylab="grain weight")
```

---

Sunflower\_Phomopsis     *Phomopsis stem canker observations for Sunflower*

---

**Description**

This dataset contains fraction intercepted photosynthetically active radiation (IPAR) and corresponding percent of girdling lesions at harvest (pclesions) for 43 fields. Phomopsis stem canker is a worldwide fungal disease of sunflower, which causes stem girdling lesions and a consequent reduction in yield. One wants to decide if the number of girdling lesions at harvest in the absence of early treatment will exceed 15 This data frame consist of a sample of fields with values for IPAR (ipar) and for the percent of girdling lesions at harvest (pclesions).

**Usage**

Sunflower\_Phomopsis

**Format**

a RangedData instance, 1 row per observation.

**Source**

Debaeke, P., & Estragnat, A. (2009). Crop canopy indicators for the early prediction of Phomopsis stem canker (*Diaporthe helianthi*) in sunflower. *Crop Protection*, 28(9), 792-801. doi:10.1016/j.cropro.2009.04.011

---

threshold.measures     *Computation of threshold.measures*

---

**Description**

Computation of threshold.measures

**Usage**

```
threshold.measures(Yobs, Ypred, p, d, units = "")
```

**Arguments**

Yobs : observed values  
 Ypred : prediction values from the model  
 p : TO COMPLETE  
 d : TO COMPLETE  
 units : units

**Value**

data.frame with the different evaluation criteria

**Examples**

```

# observed and simulated values
obs<-c(78,110,92,75,110,108,113,155,150)
sim<-c(126,126,126,105,105,105,147,147,147)
threshold.measures(obs,sim,80,1.0)

```

---

verhulst.model	<i>The Verhulst (logistic) model - calculate daily values over designated time period</i>
----------------	---

---

**Description**

The Verhulst (logistic) model - calculate daily values over designated time period

**Usage**

```
verhulst.model(a, k, Y0, duration)
```

**Arguments**

a : growth rate  
 k : capacity  
 Y0 : initial condition  
 duration : duration of simulation

**Value**

data.frame with daily Y

**See Also**

[verhulst.update](#) for the update function of the Verhulst model.

**Examples**

```
plot(verhulst.model(0.08,100,1,100), type="l", ylim=c(0,115),
     xlab="day", ylab="Y, population density",lwd=2)
```

---

verhulst.update	<i>The Verhulst (logistic) model - calculate change for one day</i>
-----------------	---

---

**Description**

The Verhulst (logistic) model - calculate change for one day

**Usage**

```
verhulst.update(Y, a, k)
```

**Arguments**

Y	: state variable $Y(t=\text{day})$
a	: growth rate
k	: capacity

**Value**

state variable at  $Y(t=\text{day}+1)$

**See Also**

[verhulst.model](#) for the integration loop function of the Verhulst model.

---

watbal.define.param	<i>Define values of the parameters for the WaterBalance model</i>
---------------------	---

---

**Description**

Define values of the parameters for the WaterBalance model

**Usage**

```
watbal.define.param()
```

**Value**

matrix with parameter values (nominal, binf, bsup)

---

watbal.model	<i>WaterBalance model - calculate soil water over designated time period</i>
--------------	--

---

### Description

WaterBalance model - calculate soil water over designated time period

### Usage

```
watbal.model(param, weather, WP, FC, WAT0 = NA)
```

### Arguments

param	: a vector of parameters
weather	: weather data.frame for one single year
WP	: Water content at wilting Point (cm <sup>3</sup> .cm <sup>-3</sup> )
FC	: Water content at field capacity (cm <sup>3</sup> .cm <sup>-3</sup> )
WAT0	: Initial Water content (mm). If NA WAT0=z*FC

### Value

data.frame with daily RAIN, ETR, Water at the beginning of the day (absolute : WAT, mm and relative value : WATp, -)

---

watbal.model.arid	<i>WaterBalance model - Variant with another order of calculation and ARID index</i>
-------------------	--

---

### Description

WaterBalance model - Variant with another order of calculation and ARID index

### Usage

```
watbal.model.arid(WHC, MUF, DC, z, CN, weather, WP, FC, WAT0 = NA)
```

**Arguments**

WHC	: Water Holding Capacity of the soil (cm <sup>3</sup> cm <sup>-3</sup> )
MUF	: Water Uptake coefficient (mm <sup>3</sup> mm <sup>-3</sup> )
DC	: Drainage coefficient (mm <sup>3</sup> mm <sup>-3</sup> )
z	: root zone depth (mm)
CN	: Runoff curve number
weather	: weather data.frame for one single year
WP	: Water content at wilting Point (cm <sup>3</sup> .cm <sup>-3</sup> )
FC	: Water content at field capacity (cm <sup>3</sup> .cm <sup>-3</sup> )
WAT0	: Initial Water content (mm). If NA WAT0=z*FC

**Value**

data.frame with daily RAIN, ETR, Water at the beginning of the day (absolute : WAT, mm and relative value : WATp, -)

---

watbal.simobsdata	<i>Soil water content measurements and associated simulations with WaterBalance model</i>
-------------------	---

---

**Description**

Data of soil water content from Luc Champolivier (CETIOM), in En Crambade (31, France), on canola without irrigation in 2008. sonde Diviner 2000 (from Sentek Pty Ltd) Simulation are from watbal.model, with an initial water content estimated from measurement with Diviner 2000.

**Usage**

```
data(watbal.simobsdata)
```

**Format**

a RangedData instance, 1 row per day : Weather : day / RAIN / ETr / simulation : WAT / WATp / ARID observation : t1\_WATp\_0\_40cm / t2\_WATp\_0\_40cm / t3\_WATp\_0\_40cm / WATp\_SF.mean / WATp\_SF.var

**Source**

CETIOM, Luc Champolivier (2008).

---

watbal.update	<i>WaterBalance model - calculate change in soil water for one day</i>
---------------	--

---

**Description**

WaterBalance model - calculate change in soil water for one day

**Usage**

```
watbal.update(WAT0, RAIN, ETr, param, WP, FC)
```

**Arguments**

WAT0	: Water at the beginning of the day (mm).
RAIN	: Rainfall of day (mm)
ETr	: Evapotranspiration of day (mm)
param	: a vector of parameters
WP	: Water content at wilting Point (cm <sup>3</sup> .cm <sup>-3</sup> )
FC	: Water content at field capacity (cm <sup>3</sup> .cm <sup>-3</sup> )

**Value**

WAT1 : Water at the beginning of the day+1 (mm).

---

watbal.weather	<i>Read weather data for the WaterBalance model (West of France Weather)</i>
----------------	--

---

**Description**

Read weather data for the WaterBalance model (West of France Weather)

**Usage**

```
watbal.weather(working.year = NA, working.site = NA, weather = NA)
```

**Arguments**

working.year	year for the subset of weather data (default=NA : all the year)
working.site	site for the subset of weather data (default=NA : all the site)
weather	weather table with different year and site

**Value**

data.frame with daily weather data for one or several site(s) and for one or several year(s)

---

weather\_EuropeEU

*Weather Series for Europe (EU) from NASA POWER Agroclimatology*

---

## Description

A contemporary daily climate dataset for Europe, covering the period from 1st January 2001 to 31 December 2010 (10 complete years of data). This dataset focuses on regions of Europe with an elevation less than 500 meters.

The dataset was extracted from the NASA Langley Research Center POWER Project, which provides agroclimatology datasets (Chandler et al., 2004). It was funded through the NASA Earth Science Directorate Applied Science Program.

This climate dataset contains daily estimates of:

- Precipitation
- Mean, minimum, and maximum temperature
- Relative humidity
- Dew point
- Solar radiation
- Wind speed

The data has global coverage at one-degree resolution (approximately 111 km at the equator).

The NASA POWER agroclimatology data are derived from multiple sources:

- Solar radiation: satellite observations
- Meteorological data: Goddard Earth Observing System global assimilation model version 4 (GEOS-4)
- Precipitation: Global Precipitation Climate Project and Tropical Rainfall Measurement Mission

A full description is available at <https://power.larc.nasa.gov>.

Elevation (Altitude) was retrieved from the Aster Global Digital Elevation Model using the web-service [api.geonames.org/astergdem](http://api.geonames.org/astergdem). Samples are approximately 30m x 30m, covering latitudes between 83 degrees N and 65 degrees S. Results provide a single number for elevation in meters (Aster GDEM). Ocean areas are masked as "no data" and assigned a value of -9999.

## Usage

```
data(weather_EuropeEU)
```

**Format**

A RangedData instance with 1 row per day and the following variables:

**SRAD** Daily Insolation Incident On A Horizontal Surface (MJ/m2/day)

**T2M** Average Air Temperature At 2 m Above The Surface Of The Earth (degrees C)

**TMIN** Minimum Air Temperature At 2 m Above The Surface Of The Earth (degrees C)

**TMAX** Maximum Air Temperature At 2 m Above The Surface Of The Earth (degrees C)

**RH2M** Relative Humidity At 2 m (percent)

**TDEW** Dew/Frost Point Temperature At 2 m (degrees C)

**RAIN** Average Precipitation (mm/day)

**WIND** Wind Speed At 10 m Above The Surface Of The Earth (m/s)

**Source**

<https://power.larc.nasa.gov/> <https://asterweb.jpl.nasa.gov/gdem.asp> <https://www.geonames.org/about.html>

**Examples**

```
data(weather_EuropeEU)
head(weather_EuropeEU)
```

---

weather_FranceWest	<i>Weather series for western France from NASA POWER agroclimatology</i>
--------------------	--

---

**Description**

This contemporary daily climate dataset for West of France covers the period 1st January 1984 to 31 December 2011. The precipitation data is limited to the period Jan-1997 Aug-2009, thus only 12 complete years of data were available for analysis involving precipitation(1997 to 2009). It cover main part of West part of France defined as a rectangle. The dataset was extracted from the NASA Langley Research Center POWER Project which provide agroclimatology dataset (Chandler et al., 2004). It was funded through the NASA Earth Science Directorate Applied Science Program This climate dataset contains daily estimates of precipitation, mean, minimum and maximum temperature, relative humidity, dew point, solar radiation and wind speed with global coverage at one degree resolution (approximately 111 km at the equator). The NASA POWER agroclimatology data are derived from various sources: solar radiation from satellite observations, meteorological data from the Goddard Earth Observing System global assimilation model version 4 (GEOS-4), and precipitation from the Global Precipitation Climate Project and Topical Rainfall Measurement Mission. A full description can be found at <https://power.larc.nasa.gov>

**Usage**

```
data(weather_FranceWest)
```

**Format**

a RangedData instance, 1 row per day.

**Source**

<https://power.larc.nasa.gov/>

---

weather\_GNS

*Weather series for Gainesville (FL, USA) years 1982 and 1983*

---

**Description**

This daily climate dataset for Gainesville (FL, USA) years 1982 and 1983 covers the period 1st January 1982 to 31 December 1983 with 2 complete years of data with precipitation. The dataset was provide by JJW Jones and the university of Florida to run simulation of the publication of Muchow et al. (1990). This climate dataset contains daily estimates of precipitation (RAIN), minimum (Tmin) and maximum temperature (Tmax), solar radiation (I), photosynthically active radiation (PAR)

**Usage**

data(weather\_GNS)

**Format**

a RangedData instance, 1 row per day.

**Source**

University of Florida

---

weather\_SouthAsia

*Weather series for southern Asia from NASA POWER agroclimatology*

---

**Description**

This contemporary daily climate dataset for South Asia covers the period 1st January 1997 to 31 December 2008 with 12 complete years of data with precipitation. It cover a part of South Asia (North-East of India, Bangladesh, Myanmar, Neapal) with an elevation less than 2500m. The dataset was extracted from the NASA Langley Research Center POWER Project which provide agroclimatology dataset (Chandler et al., 2004). It was funded through the NASA Earth Science Directorate Applied Science Program This climate dataset contains daily estimates of precipitation, mean, minimum and maximum temperature, relative humidity, dew point, solar radiation and wind speed with global coverage at one degree resolution (approximately 111 km at the equator). The NASA POWER agroclimatology data are derived from various sources: solar radiation from satellite observations, meteorological data from the Goddard Earth Observing System

global assimilation model version 4 (GEOS-4), and precipitation from the Global Precipitation Climate Project and Topical Rainfall Measurement Mission. A full description can be found at <https://power.larc.nasa.gov/> Elevation (Altitude) were retrieve from Aster Global Digital Elevation Model by using the Webservice [api.geonames.org/astergdem?](https://api.geonames.org/astergdem?) Sample are: ca 30m x 30m, between 83N and 65S latitude. Result : a single number giving the elevation in meters according to aster gdem, ocean areas have been masked as "no data" and have been assigned a value of -9999 Example <https://api.geonames.org/astergdem?lat=50.01&lng=10.2&username=demo>

**Usage**

```
data(weather_SouthAsia)
```

**Format**

a RangedData instance, 1 row per day.

**Source**

<https://power.larc.nasa.gov/> and <https://asterweb.jpl.nasa.gov/gdem.asp> and <https://www.geonames.org/about.html>

---

weed.define.param	<i>Define parameter values of the Weed model</i>
-------------------	--

---

**Description**

Define parameter values of the Weed model

**Usage**

```
weed.define.param()
```

**Value**

matrix with parameter values (nominal, binf, bsup)

---

weed.model	<i>The Weed model - calculate daily values over designated time period</i>
------------	--

---

**Description**

The Weed model - calculate daily values over designated time period

**Usage**

```
weed.model(param, weed.deci)
```

**Arguments**

param : vector of the 16 parameters  
 weed.deci : decisions table for Soil, Crop et Herbicide

**Value**

data.frame with annual values of yield

---

weed.simule	<i>Wrapper function to run the Weed model multiple times (for multiple sets of inputs)</i>
-------------	--

---

**Description**

Wrapper function to run the Weed model multiple times (for multiple sets of inputs)

**Usage**

```
weed.simule(X, weed.deci)
```

**Arguments**

X : parameter matrix  
 weed.deci : decisions table for Soil, Crop et Herbicide

**Value**

matrix with Yield for year 3 for each parameter vector

---

weed.update                      *The Weed model - calculate change for one year*

---

### Description

The Weed model - calculate change for one year

### Usage

weed.update(d, S, SSBa, DSBa, Soil, Crop, Herb, param)

### Arguments

d                                 : weed density at seed emergence (plants/m2) - value for year  
 S                                 : seed production per m2 - value for year  
 SSBa                            : surface seedbank after tillage (grains/m2) - value for year  
 DSBa                            : deep seedbank after tillage (grains/m2) - value for year  
 Soil                             : value for soil decision (1 or 0)  
 Crop                             : value for crop decision (1 or 0)  
 Herb                             : value for herbicide treatment decision (1 or 0)  
 param                          : vector of the 16 parameters

### Value

a vector with values of state variables for year+1

---

WheatYieldGreece                *National Wheat Yield evolution for Greece from FAO*

---

### Description

Wheat yield time series data in Greece from 1961 to 2010 yield.

### Usage

WheatYieldGreece

### Format

a RangedData instance, 1 row per measurement. Year, Yield : Wheat Yield (hectogram/hectare = 0.0001 ton/hectare)

### Source

Food And Agricultural Organization of United Nations (FAO), FAOSTAT database, <http://faostat.fao.org>

---

 Wheat\_GPC

*Grain Protein Contents in Wheat Grains*


---

### Description

This dataset contains data for 43 plots. The column GPC corresponds to measured data, GPC.model1 and GPC.model2 correspond to the results obtained by two models. The other columns contain other information for each plot (technical practices) : number, tillage, max\_water, preceding\_crop, sow\_date, Nendwinter, Nfertilizer, SPAD, NNI, Yield.

### Usage

Wheat\_GPC

### Format

a RangedData instance, 1 row per plot.

### Source

Barbottin et al. 2008

---

 zakoks.original.model *Classical SEIR model for plant diseases from Zadoks (1971)*


---

### Description

**Model description.** This model is a classical SEIR model for plant disease. It was written from its description included in the original publication of Zadoks (1971)

### Usage

```
zakoks.original.model(  
  nlpd = 4 * 10,  
  nipd = 1 * 10,  
  dmfr = 16,  
  SITE0 = 5 * 10^9,  
  weather,  
  sdate = 145,  
  ldate = 145 + 50,  
  XLAT0 = 1  
)
```

**Arguments**

nlpd	:	latent period (in degree.day) - default value for Puccinia triticina : ~10 days at 20degC
nipd	:	infectious period (in degree.day) - default value for Puccinia triticina : ~20 days at 20degC
dmfr	:	daily multiplication factor - default value number of effective spores produced by lesion
SITE0	:	
weather	:	weather data.frame for one single year
sdate	:	starting date
ldate	:	ending date
XLAT0	:	

**Details**

This model is a classical SEIR model proposed by Zadoks (1971) to simulate epidemics of diseases of crops. It is a Susceptible-Exposed-Infectious-Removed (SEIR) model. This simple model of an epidemic is based on the epidemiological concepts "latent period", "infectious period", and "multiplication factor". The crop is considered to consist of a large but finite number of infectious sites. The physical dimensions of an infectious site roughly coincide with the reproductive unit of the parasite studied. Different pathosystems (with different infectious site definitions) can be considered with this model. A full description, is available in the original paper: The model has four essential state variables representing the number of sites in each state XVAC for vacant (healthy) sites, XLAT for latent site, XINF for infectant sites and XCTR for the cumulative total of removal (post infectious) sites. Two supplementary variables based on the state variables are used defined as XTO1 = XLAT+XINF+XCTR and XSEV = XINF+XCTR. Fluxes or rates between the state variables are defined as rocc for occupation, rapp for apparition and rrem for removal. The model has a time step of one day (dt=1). The system modeled is one hectare of a wheat crop.

**Value**

list with a data.frame with daily day, DACE, XVAC, XLAT, XINF, XCTR, XTO1, XSEV= XSEV, severity and a vector of parameter value (nlpd, nipd, dmfr, SITE0).

**Source**

Script written from equation described in Zadoks, J.C. 1971. Systems Analysis and the Dynamics of Epidemics. *Phytopathology*. 61:441-598.

**See Also**

[epirice.model](#)

**Examples**

```
weather=subset(weather_FranceWest, WEYR==1997 & idsite==39)
out=zakoks.original.model(nlpd=4*10,nipd=1*10,dmfr=16,SITE0 = 5*10^9,
weather, sdate = 145, ldate = 145+50 , XLAT0=1)
plot(out$sim$DACE,out$sim$severity, type="l")
```

# Index

AICf, 3

Bean, 4

carbonsoil.model, 5  
carbonsoil.update, 6  
carcass.define.param, 6  
carcass.EMI.model, 7  
carcass.EMI.model2, 9  
carcass.EMI.multi, 9  
carcass.EMI.simule, 10  
carcass.model, 10  
carcass\_data, 12  
carrot.weevil.model, 12  
chicks\_data, 13  
cotton.model, 14

epirice.define.param, 15  
epirice.model, 16, 17, 65  
epirice.multi.simule, 16, 16  
epirice.weather, 17  
evaluation.criteria, 17  
exponential.model, 18  
exponential.model.bis, 19  
exponential.model.ie, 19

goodness.of.fit, 20  
graph\_epid, 21  
graph\_epid\_s, 21

lactation.calf.model, 22  
lactation.calf.model2, 23  
lactation.calf.simule, 24  
lactation.define.param, 24  
lactation.machine.model, 25  
lactation.machine.model2, 26

magarey.define.param, 27  
magarey.model, 28  
magarey.model2, 28  
magarey.simule, 29

maize.data\_EuropeEU, 29  
maize.data\_MetaModelling, 30  
maize.define.param, 30, 32  
maize.model, 30, 31  
maize.model2, 32, 32  
maize.muchow.graph, 33, 34, 42–44  
maize.muchow.model, 33, 33, 42–44  
maize.multisy, 32–34, 35, 42–44  
maize.multisy240, 35  
maize.RUEtemp, 36  
maize.simule, 32, 36  
maize.simule240, 32, 37  
maize.simule\_multisy240, 32, 38  
maize.weather, 39  
maize\_cir.model, 39  
maize\_cir\_rue.model, 40  
maize\_cir\_rue\_ear.model, 41  
mm.A.fct, 33, 34, 42, 43, 44  
mm.FAS.fct, 33, 34, 42, 42, 43, 44  
mm.HI.fct, 33, 34, 42, 43, 43, 44  
mm.LN.fct, 33, 34, 42, 43, 44

param.rtriangle, 44  
param.runif, 45  
population.age.matrix.model, 45  
population.age.model, 47  
population.age.model.ode, 48  
predator.prey.model, 49

q.arg.fast.runif, 50

seedweight.data, 51  
seedweight.model, 51  
Sunflower\_Phomopsis, 52

threshold.measures, 52

verhulst.model, 53, 54  
verhulst.update, 18–20, 53, 54

watbal.define.param, 54

watbal.model, [55](#)  
watbal.model.arid, [55](#)  
watbal.simobsdata, [56](#)  
watbal.update, [57](#)  
watbal.weather, [57](#)  
weather\_EuropeEU, [58](#)  
weather\_FranceWest, [30](#), [59](#)  
weather\_GNS, [60](#)  
weather\_SouthAsia, [60](#)  
weed.define.param, [61](#)  
weed.model, [62](#)  
weed.simule, [62](#)  
weed.update, [63](#)  
Wheat\_GPC, [64](#)  
WheatYieldGreece, [63](#)  
  
zakoks.original.model, [21](#), [22](#), [64](#)