

Package ‘photosynthesisLRC’

May 9, 2026

Title Nonlinear Least Squares Models for Photosynthetic Light Response

Version 1.0.6

Description Provides functions for modeling, comparing, and visualizing photosynthetic light response curves using established mechanistic and empirical models like the rectangular hyperbola Michaelis-Menton based models ((eq1 (Baly (1935) <[doi:10.1098/rspb.1935.0026](https://doi.org/10.1098/rspb.1935.0026)>)) (eq2 (Kaipainenn (2009) <[doi:10.1134/S1021443709040025](https://doi.org/10.1134/S1021443709040025)>)) (eq3 (Smith (1936) <[doi:10.1073/pnas.22.8.504](https://doi.org/10.1073/pnas.22.8.504)>))), hyperbolic tangent based models ((eq4 (Jassby & Platt (1976) <[doi:10.4319/LO.1976.21.4.0540](https://doi.org/10.4319/LO.1976.21.4.0540)>)) (eq5 (Abe et al. (2009) <[doi:10.1111/j.1444-2906.2008.01619.x](https://doi.org/10.1111/j.1444-2906.2008.01619.x)>))), the non-rectangular hyperbola model (eq6 (Prioul & Chartier (1977) <[doi:10.1093/oxfordjournals.aob.a085354](https://doi.org/10.1093/oxfordjournals.aob.a085354)>)), exponential based models ((eq8 (Webb et al. (1974) <[doi:10.1007/BF00345747](https://doi.org/10.1007/BF00345747)>)), (eq9 (Prado & de Moraes (1997) <[doi:10.1007/BF02982542](https://doi.org/10.1007/BF02982542)>)) nally the Ye model (eq11 (Ye (2007) <[doi:10.1007/s11099-007-0110-5](https://doi.org/10.1007/s11099-007-0110-5)>)). Each of these nonlinear least squares models are commonly used to express photosynthetic response under changing light conditions and has been well supported in the literature, but distinctions in each mathematical model represent moderately different assumptions about physiology and trait relationships which ultimately produce different calculated functional trait values. These models were all thoughtfully discussed and curated by Lobo et al. (2013) <[doi:10.1007/s11099-013-0045-y](https://doi.org/10.1007/s11099-013-0045-y)> to express the importance of selecting an appropriate model for analysis, and methods were established in Davis et al. (in review) to evaluate the impact of analytical choice in phylogenetic analysis of the function-valued traits. Gas exchange data on 28 wild sunflower species from Davis et al. are included as an example data set here.

Maintainer Rebekah Davis <rebekah.davis.evoecophys@gmail.com>

Depends R (>= 4.2), dplyr (>= 1.1.4), tidyr (>= 1.3.1), stats (>= 4.2.3), graphics (>= 4.2.3)

LazyData true

URL <https://github.com/heliotropichuman/photosynthesisLRC>

License MIT + file LICENSE

Encoding UTF-8

RoxygenNote 7.3.2

NeedsCompilation no

Author Rebekah Davis [aut, cre],
Eric Goolsby [aut]

Repository CRAN

Date/Publication 2025-02-25 17:50:07 UTC

Contents

eq1	2
eq11	4
eq2	6
eq3	7
eq4	9
eq5	11
eq6	13
eq8	15
eq9	16
nls_results	18
plot_eq	21
sunflowers	22

Index **25**

eq1	<i>Calculate Photosynthetic Rates Using a Nonlinear Model EQ1</i>
-----	---

Description

Uses the nonlinear least squares Michaelis-Menton model equation 1 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim \frac{(\phi_{I0})(PAR_i)(P_{gmax})}{(\phi_{I0} * PAR_i + P_{gmax})} - R_d$ The function will return predicted values, calculated quantities, or both.

Usage

```
eq1(pars = c(Pgmax = 19.5, phi_I0 = .0899, Rd = 1.8),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])
```

Arguments

pars	A named vector of parameters. Default values are Pgmax = 19.5, phi_I0 = 0.0899, and Rd = 1.8. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. These values may be changed to fall within the minimum and maximum parameter values of your study system.
data	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the measured photosynthetic rate.
PARi	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.

return Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters P_{gmax} , ϕ_{I0} , and R_d by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: P_{gmax} , P_{max} , I_{comp} , ϕ_{I0} (quantum yield calculated at I_0), $\phi_{I_{comp}}$ (quantum yield calculated at I_{comp}), $\phi_{I_0, I_{comp}}$ (quantum yield calculated by the range of values between I_0 and I_{comp}), $\phi_{I_{comp}, I_{200}}$ (quantum yield calculated by the range between I_{comp} and I_{200}), R_d (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), $P_{I_{max}}$ (assimilation value at maximum light), I_{sat_x} , $x = .25, .50, .75, .85, .90, .95$ (light saturation at x percent of P_{max}), I_x , $x = .25, .50, .75, .85, .90, .95$ (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli

Baly, E.C.C. 1935 The kinetics of photosynthesis. Proc. R. Soc. Lond. B. 117: 218-239.

Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARI = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq1(pars = c(Pgmax = 20, phi_I0 = 0.09, Rd = 2),
  PARI = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate
# linear parameters
```

```

result <- eq1(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq1(data = example_data, return = "calc")
print(calculated_quantities)

```

eq11

Calculate Photosynthetic Rates Using a Nonlinear Model EQ11

Description

Uses the nonlinear least squares Ye model equation 11 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim \phi_{I0} I_{comp} \frac{(1 - \beta(PAR_i))}{(1 + \gamma(PAR_i))} (PAR_i - I_{comp})$ The function will return predicted values, calculated quantities, or both.

Usage

```

eq11(pars = c(phi_I0_Icomp = .0756, beta = .0000432, gamma = .0039, Icomp = 22.6),
     data,
     PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
     return = c("predict", "calc", "all")[1])

```

Arguments

pars	A named vector of parameters. Default values are $\phi_{I0_Icomp} = .0756$, $\beta = .0000432$, $\gamma = .0039$, $I_{comp} = 22.6$. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. The empirical values β and γ range between 0 and 1, and are not explicitly described by Ye (2007) but are independent coefficients of I implemented to incorporate a more dynamic response to light. All of these values may be changed to fall within the minimum and maximum parameter values of your study system.
data	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
PARi	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
return	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters ϕ_{I_0} , I_{comp} , and empirical parameters β and γ by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: P_{gmax} , P_{max} , I_{comp} , ϕ_{I_0} (quantum yield calculated at I_0), $\phi_{I_{comp}}$ (quantum yield calculated at I_{comp}), $\phi_{I_0, I_{comp}}$ (quantum yield calculated by the range of values between I_0 and I_{comp}), $\phi_{I_{comp}, I_{200}}$ (quantum yield calculated by the range between I_{comp} and I_{200}), R_d (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), $P_{I_{max}}$ (assimilation value at maximum light), I_{sat_x} , $x = .25, .50, .75, .85, .90, .95$ (light saturation at x percent of P_{max}), I_x , $x = .25, .50, .75, .85, .90, .95$ (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

- Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli
- Ye, Z.-P. 2007 A new model for relationship between irradiance and the rate of photosynthesis in *Oryza s
- Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
data(sunflowers)
example_data <- sunflowers |> filter(SampleID==SampleID[1])
# Predict photosynthetic rates given the parameters
predicted_values <- eq11(return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate linear parameters
result <- eq11(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq11(data = example_data, return = "calc")
print(calculated_quantities)
```

eq2

*Calculate Photosynthetic Rates Using a Nonlinear Model EQ2***Description**

Uses the nonlinear least squares Michaelis-Menton model equation 2 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim ((P_{gmax} * PAR_i) / (PAR_i + I_{50})) - R_d$ The function will return predicted values, calculated quantities, or both.

Usage

```
eq2(pars = c(Pgmax = 19.5, I50 = 216.4, Rd = 1.8),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])
```

Arguments

pars	A named vector of parameters. Default values are $P_{gmax} = 19.5$, $I_{50} = 216.4$, and $R_d = 1.8$. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. These values may be changed to fall within the minimum and maximum parameter values of your study system.
data	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
PARi	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
return	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters P_{gmax} , assimilation at half the maximum assimilation rate (I_{50}), and R_d by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.

- "calc": A named vector of calculated quantities: Pgmax, Pmax, Icomp, phi_I0 (quantum yield calculated at I0), phi_Icomp (quantum yield calculated at Icomp), phi_I0_Icomp (quantum yield calculated by the range of values between I0 and Icomp), phi_Icomp_I200 (quantum yield calculated by the range between Icomp and I200), Rd (dark respiration), Imax (Imax calculated), Imax_obs (Imax observed), P_Imax (assimilation value at maximum light), Isat_x, x = .25, .50, .75, .85, .90, .95 (light saturation at x percent of Pmax), Ix, x = .25, .50, .75, .85, .90, .95 (light intensity at x percent of Pmax)
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

- Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli
- Kaipiainen, E.L. 2009 Parameters of photosynthesis light curve in *Salix dasyclados* and their changes d
- Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq2(pars = c(Pgmax = 20, I50 = 216.4, Rd = 2),
  PARi = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate
# linear parameters
result <- eq2(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq2(data = example_data, return = "calc")
print(calculated_quantities)
```

Description

Uses the nonlinear least squares Michaelis-Menton model equation 3 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim ((\phi_{I0})(PAR_i)(P_{gmax}) / (((P_{gmax}^2) + (\phi_{I0}^2) * (PAR_i^2))^{0.5}) - R_d$ The function will return predicted values, calculated quantities, or both.

Usage

```
eq3(pars = c(Pgmax = 19.5, phi_I0 = .0493, Rd = 1.8),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])
```

Arguments

pars	A named vector of parameters. Default values are Pgmax = 19.5, phi_I0 = .0493, and Rd = 1.8. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. These values may be changed to fall within the minimum and maximum parameter values of your study system.
data	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
PARi	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
return	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters Pgmax, phi_I0, and Rd by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (Rd), light compensation point (Icomp), maximum photosynthetic rate (Pmax), and curve derived parameters (Ix) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: Pgmax, Pmax, Icomp, phi_I0 (quantum yield calculated at I0), phi_Icomp (quantum yield calculated at Icomp), phi_I0_Icomp (quantum yield calculated by the range of values between I0 and Icomp), phi_Icomp_I200 (quantum yield calculated by the range between Icomp and I200), Rd (dark respiration), Imax (Imax calculated), Imax_obs (Imax observed), P_Imax (assimilation value at maximum light), Isat_x, x = .25, .50, .75, .85, .90, .95 (light saturation at x percent of Pmax), Ix, x = .25, .50, .75, .85, .90, .95 (light intensity at x percent of Pmax)
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

- Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli
- Smith, E. L. 1936 Photosynthesis in relation to light and carbon dioxide. PNAS 22: 504-511.
- Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq3(pars = c(Pgmax = 20, phi_I0 = .0493, Rd = 2),
  PARi = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate
# linear parameters
result <- eq3(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq3(data = example_data, return = "calc")
print(calculated_quantities)
```

 eq4

Calculate Photosynthetic Rates Using a Nonlinear Model EQ4

Description

Uses the nonlinear least squares hyperbolic tangent model equation 4 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim P_{gmax} * \tanh((\phi_{I0}) PAR_i / P_{gmax}) - R_d$ The function will return predicted values, calculated quantities, or both.

Usage

```
eq4(pars = c(Pgmax = 19.5, phi_I0 = .0493, Rd = 1.8),
  data,
  PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  return = c("predict", "calc", "all")[1])
```

Arguments

<code>pars</code>	A named vector of parameters. Default values are $P_{gmax} = 19.5$, $\phi_{I0} = 0.0493$, and $R_d = 1.8$. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. These values may be changed to fall within the minimum and maximum parameter values of your study system.
<code>data</code>	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
<code>PARi</code>	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
<code>return</code>	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters P_{gmax} , ϕ_{I0} , and R_d by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: P_{gmax} , P_{max} , I_{comp} , ϕ_{I0} (quantum yield calculated at I_0), $\phi_{I_{comp}}$ (quantum yield calculated at I_{comp}), $\phi_{I_0 I_{comp}}$ (quantum yield calculated by the range of values between I_0 and I_{comp}), $\phi_{I_{comp} I_{200}}$ (quantum yield calculated by the range between I_{comp} and I_{200}), R_d (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), $P_{I_{max}}$ (assimilation value at maximum light), I_{sat_x} , $x = .25, .50, .75, .85, .90, .95$ (light saturation at x percent of P_{max}), I_x , $x = .25, .50, .75, .85, .90, .95$ (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

- Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli
- Abe, M., K. Yokota, A. Kurashima, M. Maegawa 2009 High water temperature tolerance in photosynthetic act
- Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
```

```

    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
  )

# Predict photosynthetic rates given the parameters
predicted_values <- eq4(pars = c(Pgmax = 20, phi_I0 = 0.09, Rd = 2),
                       PARi = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate
# linear parameters
result <- eq4(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit)  # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq4(data = example_data, return = "calc")
print(calculated_quantities)

```

eq5

Calculate Photosynthetic Rates Using a Nonlinear Model EQ5

Description

Uses the nonlinear least squares hyperbolic tangent model equation 5 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim Pgmax * \tanh(PARi / Isat) - Rd$ The function will return predicted values, calculated quantities, or both.

Usage

```

eq5(pars = c(Pgmax = 15.5, Isat = 359.2, Rd = .9),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])

```

Arguments

pars	A named vector of parameters. Default values are Pgmax = 15.5, Isat = 359.2, and Rd = 0.9. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. These values may be changed to fall within the minimum and maximum parameter values of your study system.
data	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
PARi	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.

return Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters P_{gmax} , I_{sat} , and R_d by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: P_{gmax} , P_{max} , I_{comp} , ϕ_{I0} (quantum yield calculated at I_0), $\phi_{I_{comp}}$ (quantum yield calculated at I_{comp}), $\phi_{I_0-I_{comp}}$ (quantum yield calculated by the range of values between I_0 and I_{comp}), $\phi_{I_{comp}-I_{200}}$ (quantum yield calculated by the range between I_{comp} and I_{200}), R_d (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), $P_{I_{max}}$ (assimilation value at maximum light), I_{sat_x} , $x = .25, .50, .75, .85, .90, .95$ (light saturation at x percent of P_{max}), I_x , $x = .25, .50, .75, .85, .90, .95$ (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

- Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli
- Jassby, A.D. and T. Platt 1976 Mathematical formulation of the relationship between photosynthesis and I
- Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARI = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq5(pars = c(Pgmax = 15.5, Isat = 360, Rd = .9),
  PARI = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate linear parameters
result <- eq5(data = example_data, return = "all")
```

```

print(result$calc) # View calculated quantities
print(result$fit)  # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq5(data = example_data, return = "calc")
print(calculated_quantities)

```

eq6

*Calculate Photosynthetic Rates Using a Nonlinear Model EQ6***Description**

Uses the nonlinear least squares non-rectangular hyperbola model equation 6 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim \frac{((\text{PAR}_i)(\phi_{I0}) + P_{gmax}) - (((\phi_{I0})(\text{PAR}_i) + P_{gmax})^2) - (4(\phi_{I0})(P_{gmax})(\theta)(\text{PAR}_i)^{0.5})}{(2 * \text{exp}(\text{Rd}))}$ The function will return predicted values, calculated quantities, or both.

Usage

```

eq6(pars = c(Pgmax = 15.5, phi_I0 = .0493, theta = .433, Rd = .9),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])

```

Arguments

<code>pars</code>	A named vector of parameters. Default values are <code>Pgmax= 15.5</code> , <code>phi_I0= 0.0493</code> , <code>theta= 0.433</code> , and <code>Rd= 0.9</code> . These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. The empirical coefficient <code>theta</code> falls between 0 and 1, and represents the various resistances faced by CO ₂ as it diffuses through the leaf mesophyll and is eventually bound by carboxylation enzymes. All of these values may be changed to fall within the minimum and maximum parameter values of your study system.
<code>data</code>	A data frame containing the experimental data with at least two columns: <code>'PARi'</code> for the incident light and <code>'A'</code> for the photosynthetic rate.
<code>PARi</code>	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
<code>return</code>	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters `Pgmax`, `phi_I0`, and `Rd` by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (`Rd`), light compensation point (`Icomp`), maximum photosynthetic rate (`Pmax`), and curve derived parameters (`Ix`) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: P_{gmax}, P_{max}, I_{comp}, phi_I0 (quantum yield calculated at I0), phi_I_{comp} (quantum yield calculated at I_{comp}), phi_I0_I_{comp} (quantum yield calculated by the range of values between I0 and I_{comp}), phi_I_{comp}_I200 (quantum yield calculated by the range between I_{comp} and I200), Rd (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), P_I_{max} (assimilation value at maximum light), Isat_x, x = .25, .50, .75, .85, .90, .95 (light saturation at x percent of P_{max}), Ix, x = .25, .50, .75, .85, .90, .95 (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli

Prioul, J. L., P. Chartier 1977 Partitioning of transfer and carboxylation components of intracellular r

Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq6(pars = c(Pgmax = 15.5, phi_I0 = .0493,
  theta = .433,Rd = .9),PARi = c(0, 100, 200, 400, 800),
  return= "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate linear parameters
result <- eq6(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq6(data = example_data, return = "calc")
print(calculated_quantities)
```

Description

Uses the nonlinear least squares exponential model equation 8 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim (P_{gmax} * (1 - \exp(-\phi_{I0}(PAR_i)/P_{gmax}))) - R_d$ The function will return predicted values, calculated quantities or both.

Usage

```
eq8(pars = c(Pgmax = 16.2, phi_I0 = .0597, Rd = 1.3),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])
```

Arguments

<code>pars</code>	A named vector of parameters. Default values are $P_{gmax} = 16.2$, $\phi_{I0} = .0597$, and $R_d = 1.3$. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. These values may be changed to fall within the minimum and maximum parameter values of your study system.
<code>data</code>	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
<code>PARi</code>	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
<code>return</code>	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters P_{gmax} , ϕ_{I0} , and R_d by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.

- "calc": A named vector of calculated quantities: P_{gmax}, P_{max}, I_{comp}, phi_I0 (quantum yield calculated at I0), phi_Icomp (quantum yield calculated at Icomp), phi_I0_Icomp (quantum yield calculated by the range of values between I0 and Icomp), phi_Icomp_I200 (quantum yield calculated by the range between Icomp and I200), Rd (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), P_I_{max} (assimilation value at maximum light), Isat_x, x = .25, .50, .75, .85, .90, .95 (light saturation at x percent of P_{max}), Ix, x = .25, .50, .75, .85, .90, .95 (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli

Webb, W. L., M. Newton, D. Starr 1974 Carbon dioxide exchange of *Alnus rubra*: a mathematical model. Oec

Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARI = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq8(pars = c(Pgmax = 16, phi_I0 = .0597, Rd = 1.3),
  PARI = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate linear parameters
result <- eq8(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq8(data = example_data, return = "calc")
print(calculated_quantities)
```

eq9

Calculate Photosynthetic Rates Using a Nonlinear Model EQ9

Description

Uses the nonlinear least squares exponential model equation 9 from Lobo et. al (2013) to transform measured photosynthetic data into a smoothed function-valued trait with the following function: $A \sim P_{gmax}((1 - \exp(-k \cdot (PARI - I_{comp})))) - R_d$ The function will return predicted values, calculated quantities, or both.

Usage

```
eq9(pars = c(Pgmax = 22, Icomp = 10, k = .0015, Rd = .1),
    data,
    PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
    return = c("predict", "calc", "all")[1])
```

Arguments

<code>pars</code>	A named vector of parameters. Default values are $P_{gmax} = 22$, $I_{comp} = 10$, $k = .0015$, and $R_d = .1$. These serve as initial starting parameters for the function to rapidly assess your data through an iterative process. The coefficient k relates the photosynthetic capacity of a plant to the specific leaf mass, and may increase with an increasing potential for photosynthetic activity per unit leaf mass. All initial parameter values may be changed to fall within the minimum and maximum parameter values of your study system.
<code>data</code>	A data frame containing the experimental data with at least two columns: 'PARi' for the incident light and 'A' for the photosynthetic rate.
<code>PARi</code>	A numeric vector of incident light values. Defaults to a sequence from 0 to 2500.
<code>return</code>	Character string indicating what the function should return. Options are "predict" for predicted values, "calc" for calculated quantities, and "all" for both. Defaults to "predict".

Details

The function uses the provided data to estimate the parameters P_{gmax} , I_{comp} , R_d , and k by minimizing the squared differences between observed and predicted photosynthetic rates. The model is then used to calculate a range of derived functional trait quantities such as the dark respiration rate (R_d), light compensation point (I_{comp}), maximum photosynthetic rate (P_{max}), and curve derived parameters (I_x) among other calculated quantities.

Value

Depending on the 'return' argument, the function returns:

- "predict": A numeric vector of predicted photosynthetic rates.
- "calc": A named vector of calculated quantities: P_{gmax} , P_{max} , I_{comp} , ϕ_{I0} (quantum yield calculated at I_0), $\phi_{I_{comp}}$ (quantum yield calculated at I_{comp}), $\phi_{I_0-I_{comp}}$ (quantum yield calculated by the range of values between I_0 and I_{comp}), $\phi_{I_{comp}-I_{200}}$ (quantum yield calculated by the range between I_{comp} and I_{200}), R_d (dark respiration), I_{max} (I_{max} calculated), I_{max_obs} (I_{max} observed), $P_{I_{max}}$ (assimilation value at maximum light), I_{sat_x} , $x = .25, .50, .75, .85, .90, .95$ (light saturation at x percent of P_{max}), I_x , $x = .25, .50, .75, .85, .90, .95$ (light intensity at x percent of P_{max})
- "all": A list containing both the predicted values, calculated quantities, and model fit statistics.

References

- Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourli
- Prado, C. H. B. A., J. P. A. P. V. de Moraes 1997 Photosynthetic capacity and specific leaf mass in twenty
- Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curv

Examples

```
# Example dataset
example_data <- data.frame(
  PARI = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
  A = c(1.8, 4.2, 7.5, 12.8, 16.2, 18.5, 19.3, 19.4, 19.5)
)

# Predict photosynthetic rates given the parameters
predicted_values <- eq9(pars= c(Pgmax = 20,Icomp = 20,k = .0015,Rd = .2),
  PARI = c(0, 100, 200, 400, 800), return = "predict")
print(predicted_values)

# Use experimental data to predict photosynthetic rates and estimate linear parameters
result <- eq9(data = example_data, return = "all")
print(result$calc) # View calculated quantities
print(result$fit) # View fit statistics and optimized parameters

# Get calculated quantities directly
calculated_quantities <- eq9(data = example_data, return = "calc")
print(calculated_quantities)
```

nls_results

Array of Results from all NLS Models eq1-eq11

Description

This function generates an array of results from the NLS models presented in Lobo et al. (2013) and Davis et al. (2024). The array includes all calculated values/model parameters for each NLS model (eq), predicted values of the NLS curve, and goodness of fit statistics - mean squared error (MSE) and r2 values- for each predicted curve and calculated value.

Usage

```
nls_results(
  data,
  species_col,
  sample_col,
  A_col,
  specs = NULL,
```

```

dat_wide = NULL,
PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500),
PARi_fine = seq(0,6000,by = .1),
eqs = paste("eq", c(1, 2, 3, 4, 5, 6, 8, 9, 11), sep = ""),
par_names = c("Pgmax", "Pmax", "phi_I0", "phi_I0_Icomp", "phi_Icomp",
              "phi_Icomp_200", "Rd", "Icomp", "Isat", "Isat_25",
              "Isat_50", "Isat_75", "Isat_85", "Isat_90", "Isat_95",
              "I5", "I10", "I15", "I25", "I50", "I75", "I85", "I90",
              "I95", "Imax", "Imax_obs", "P_Imax",
              "beta", "gamma", "theta", "k"))

```

Arguments

data	A data frame of measured input data with a column for each unique plant observed, a column to indicate a common identifier among each unique observation a column of light intensities at measured PARi values, and a column for measured measured carbon assimilation values.
species_col	Identify the column in your data set for the Species name, or the common identifier for each sample (Ex. H.Debilis)
sample_col	Identify the column in your data set with the Sample ID, or the unique identifier for each observation (Ex. H.Debilis_2_Oct1).
A_col	Identify the column in your data set corresponding to the measured carbon assimilation value (A).
specs	A character or factor vector to subset individual identifiers, defaults to all unique samples in the data set.
dat_wide	A wide-format data frame used to extract observed PARi and A values for each Species and Sample taken, defaults to transform dat_wide from data automatically.
PARi	A numeric vector of PARi values in which carbon assimilation was measured. Defaults to c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500).
PARi_fine	A finer numeric vector of PARi values for estimating response across a continuous light range, defaults to seq(0,6000,by = .1) for fine scale predictions.
eqs	A character vector of equation names to be fitted. Defaults to the 9 equations programmed in this package: (eq1, eq2, eq3, eq4, eq5, eq6, eq8, eq9, eq11).
par_names	A character or factor vector of parameters calculated from the NLS models to include in the array of results. Defaults to parameters calculated from all 9 models in this package unless otherwise specified.

Details

The function fits each equation from the eqs list to the subset of data corresponding to each unique sample. It then returns calculated parameters, predicted values, and observed PARi values for the given individual(s). The results are stored in an array, and the goodness-of-fit metrics (r^2 and mse) are saved in separate matrices.

Value

A list containing the following elements:

res An array of results. The array dimensions are: number of photosynthetic models (eq) tested x (number of parameters + 2 * length(PARi)) x length(inds).

fitted_curves A list of fitted curves for each equation and individual.

r2 A matrix of r2 values for each individual and equation.

mse A matrix of mse values for each individual and equation.

References

Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curve: approaches and pitfalls in phylogenetic modeling of a function-valued trait. IJPS, in review

Lobo, F. de A., M. P. de Barros, H. J. Dalmagro, .C. Dalmolin, W. E. Pereira, É.C. de Souza, G. L. Vourlitis and C. E. Rodriguez Ortiz 2013 Fitting net photosynthetic light-response curves with Microsoft Excel - a critical look at the models. Photosynthetica 51 (3): 445-456.

Examples

```
# Example dataset
data(sunflowers)
data <- sunflowers |>filter(SampleID==SampleID)

# Specify arguments
PARi = c(0, 50, 100, 250, 500, 1000, 1500, 2000, 2500)
PARifine <- seq(0, 1500, by = 1)
par_names <- c("Pmax", "Icomp", "phi_Icomp", "Rd", "I15, I25", "I85", "I95")
dat_wide <- data %>% pivot_wider(id_cols = c(Species,SampleID),
                               names_from = PARi, names_prefix = "PARi_",values_from = A)
subset <- first_five <- unique(data$SampleID[1:43])

# Process light response curve data for all model equations
my_results <- nls_results(data = data, species_col = "Species",
                        sample_col = "SampleID", A_col = "A", PARi_fine = PARifine,
                        specs = subset, dat_wide = dat_wide, par_names = par_names)

# Access the results array
result_array <- my_results$res
# Access the fitted curves
fitted_curves <- my_results$fitted_curves
# Access r2 and mse matrices
r2_matrix <- my_results$r2
mse_matrix <- my_results$mse
```

Description

This function plots the fit of a given photosynthetic light response equation for all or select species in a data set. This base R plot includes the SampleID and the equation number in the title.

Usage

```
plot_eq(eqX,
        eq_name,
        i,
        data,
        title,
        species_subset = NULL,
        A_col)
```

Arguments

eqX	A function representing the photosynthetic light response equation (e.g., eq1, eq2).
eq_name	A character string representing the name of the equation, to be included in the plot title.
i	An integer specifying the index of the species in the inds vector.
data	A data frame containing the experimental data with at least two columns: PAR _i for the incident light and A for the measured photosynthetic rate.
title	An optional character string specifying the title of the plot (defaults to title in the format i SampleID Equation X).
species_subset	An optional vector of species names from inds to be plotted. If NULL, all species in inds will be used (default is NULL).
A_col	Allows data column with assimilation measurements to be specified and defaults to A.

Details

This function takes the equation of photosynthetic light response models and fits it to the data for a given species. It then plots the observed and predicted values, highlighting specific points on the curve (such as the model curve parameters I₁₅, I₂₅, I₈₅, and I₉₅), where the number (X) is the carbon assimilation rate at X percent of the maximum assimilation in the measured data. The equation name is included in the plot title, and an optional subset of species can be selected for plotting. The function also calculates various fit statistics and adds both the original and reconstructed predictions as curves to the plot.

Value

A plot of the measured data points for the selected species (open points), with curve parameters from the fitted equation (black points), the NLS curve (red line), and the model fit (dashed blue line). It will also return the reconstructed model fit as a list.

Examples

```
# Example with eq1 and all species
# Please note, it may take more than 10 seconds to plot graphs with all species

data(sunflowers)
my_observed_data <- sunflowers
inds <- unique(my_observed_data$SampleID)

# Example with eq1 and all species

for (i in 1:length(inds)) {
  plot_eq(eq1, "eq1", i, data = my_observed_data)
}

# Example of using the function for all equations with all species or a subset of species

LRCdata <- sunflowers |> filter(SampleID==SampleID)
highlight <- c("Agrestis_1_29/10/19", "Atrorubens_3_11/11/2019", "Divaricatus_2_29/10/19",
"Gracilentus_2_3/11/2019", "Gracilentus_5_5/11/2019", "Silphiodias_1_3/11/2019")
par(mfrow = c(3, 3))

for (i in 1:length(highlight)) {
  # Add equation names to the function calls
  plot_eq(eq1, "eq1", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq2, "eq2", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq3, "eq3", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq4, "eq4", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq5, "eq5", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq6, "eq6", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq8, "eq8", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq9, "eq9", i, data = LRCdata, species_subset = highlight)
  plot_eq(eq11, "eq11", i, data = LRCdata, species_subset = highlight)
  dev.off()
}
oldpar<- par(mfrow = c(1,2))
par(oldpar)
on.exit()
```

Description

This dataset contains photosynthetic light response curves for 28 sunflower species in the genus *Helianthus*. The data was gathered in a common garden to evaluate phylogenetic change to photosynthetic response across as a function-valued trait (Davis et al. 2024).

Usage

sunflowers

Format

A data frame with 666 rows and 22 variables:

SampleID A unique identifier for each sample, a combination of Species_Replicate_Date (character).

Species The species name of the sunflower (character).

Replicate The number assigned to indicate repeated individuals of the same species (numeric).

Date The day at which the measurement was taken (character, DD:MM:YY format).

Time The time at which the measurement was taken (character, HH:MM format).

CO2r The TARGAS-1 reference CO₂ (μmol CO₂ mol⁻¹, numeric).

CO2a The TARGAS-1 ambient CO₂ reading (μmol CO₂ mol⁻¹, numeric).

H2Or Humidity reading measured by the TARGAS-1 reference during the time of measurement (mb, numeric).

H2Oa Humidity reading measured in the air by the TARGAS-1 during the time of measurement (mb, numeric).

atm Atmospheric pressured at the measurement (mb, numeric).

Flow.Supply Air flow moving through the TARGAS-1 (cc/min, numeric).

Flow.sample Air flow passing back into the TARGAS-1 from the PLC (cc/min, numeric).

PARe Photosynthetically Active Radiation levels in the environment, measured at the PLC (μmol m⁻² s⁻¹, numeric).

PARi Photosynthetically Active Radiation levels on the leaf (μmol m⁻² s⁻¹, numeric).

Tleaf Leaf temperature in the PLC during the measurement (degrees Celsius, numeric).

Trans Transpiration rate (mmol H₂O m⁻² s⁻¹, numeric).

VPD Vapor pressure deficit during the measurement (mb, numeric).

gs Stomatal conductance (mmol H₂O m⁻² s⁻¹, numeric).

A Photosynthetic rate (μmol CO₂ m⁻² s⁻¹, numeric).

Ci Concentration of leaf internal CO₂ (μmol CO₂ mol⁻¹, numeric).

Area The area of the leaf measured (cm², numeric).

Details

All plants were grown in fertilized 1 gallon pots during August- December 2019 at the University of Central Florida's Transgenic Greenhouse. Measurements were taken for two hours before and after solar noon with the Portable Photosynthetic Systems (PPS) TARGAS-1 on the most recently expanded full leaf during the juvenile stage just before flowering (V4-6) (NDSU Stages of Sunflower Development). The TARGAS-1 portable leaf cuvette (PLC) was clamped on the leaf perpendicular to the leaf margin, and photosynthetic measurements were taken at approximately 420 ppm CO₂, ambient humidity, and 500 ppm flow. Photosynthetic responses were recorded at 9 levels with an acclimation of 2-5 minutes at PARi 0, PARi 50, PARi 100, PARi 250, PARi 500, PARi 1000, PARi 2000, PARi 2500 ($\mu\text{mol s/m}^2$). The data was manually curated such that duplicate instantaneous assimilation points were removed and only one single representative sample from each replicate was used in analysis. The data set includes the Sample name, the species name, the replicate number, and 19 other variables obtained from the TARGAS-1 measurement.

Author(s)

Rebekah Davis, University of Central Florida

Source

Data collected by Rebekah Davis at the University of Central Florida.

References

Davis, R.E., C. M. Mason, E. W. Goolsby 2024 Comparative evolution of photosynthetic light response curve: approaches and pitfalls in phylogenetic modeling of a function-valued trait. IJPS, in review

Kandel, H., A. A. Schneiter, J. F. Miller, D. R. Berglund 2019 Stages of Sunflower Development. North Dakota State University Extension (<https://www.ndsu.edu/agriculture/extension/publications/stages-sunflower-development>)

Examples

```
# Load the dataset
data(sunflowers)

# View the first few rows
head(sunflowers)

# Summary statistics for photosynthetic rate (A)
summary(sunflowers$A)

# Plot the photosynthetic rate vs. PARi
plot(sunflowers$PARi, sunflowers$A,
     xlab = "PAR ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ )",
     ylab = "Photosynthetic rate ( $\mu\text{mol CO}_2 \text{ m}^{-2} \text{s}^{-1}$ )")
```

Index

* dataset

sunflowers, [22](#)

eq1, [2](#)

eq11, [4](#)

eq2, [6](#)

eq3, [7](#)

eq4, [9](#)

eq5, [11](#)

eq6, [13](#)

eq8, [15](#)

eq9, [16](#)

nls_results, [18](#)

plot_eq, [21](#)

sunflowers, [22](#)